arXiv:math/0604006v2 [math.SP] 10 Jul 2006

Schrödinger operators on zigzag graphs

Evgeny Korotyaev [*] Igor Lobanov [†]

2nd February 2008

Abstract

We consider the Schrödinger operator on zigzag graphs with a periodic potential. The spectrum of this operator consists of an absolutely continuous part (intervals separated by gaps) plus an infinite number of eigenvalues with infinite multiplicity. We describe all compactly supported eigenfunctions with the same eigenvalue. We define a Lyapunov function, which is analytic on some Riemann surface. On each sheet, the Lyapunov function has the same properties as in the scalar case, but it has branch points, which we call resonances. We prove that all resonances are real. We determine the asymptotics of the periodic and anti-periodic spectrum and of the resonances at high energy. We show that there exist two types of gaps: i) stable gaps, where the endpoints are periodic and anti-periodic eigenvalues, ii) unstable (resonance) gaps, where the endpoints are resonances (i.e., real branch points of the Lyapunov function). We obtain the following results from the inverse spectral theory: 1) we describe all finite gap potentials, 2) the mapping: potential – all eigenvalues is a real analytic isomorphism for some class of potentials.
We apply all these results to quasi-1D models of zigzag single-well carbon nanotubes.

1 Introduction and main results

Consider the Schrödinger operator $H = -\partial^2 + q$ with a periodic potential q on the zigzag graph Γ^N. The zigzag graph Γ^N is related to a zigzag nanotube T^N. We shall discuss this at the end of this section. In order to define Γ^N we introduce the so-called honeycomb lattice $\Gamma = \cup_\omega \Gamma_\omega \subset \mathbb{R}^2$, $\omega = (n, j, k) \in \mathbb{Z} \times \mathbb{J} \times \mathbb{Z}, \mathbb{J} = \{0, 1, 2\}$, where the edges Γ_ω are given by

$$\Gamma_\omega = \{x = r_\omega + te_\omega, \quad t \in [0, 1]\}, \quad e_\omega = e_j \in \mathbb{R}^2, \quad r_{(n,1,k)} = r_{(n,0,k)} + e_0,$$

$$r_{(n,0,k)} = r_{(n-1,2,k)} = n(e_0 + e_1) + k(e_1 + e_2), \quad e_0 = (0, 1),\ e_1 = \frac{1}{2}(\sqrt{3}, 1),\ e_2 = \frac{1}{2}(\sqrt{3}, -1),$$

see Fig. 1 and 2. The edges Γ_ω of length 1 form regular hexagons. Each edge Γ_ω has orientation given by $e_\omega = e_j$ with the starting point $r_\omega \in \mathbb{R}^2$. We have the coordinate

---

[*] Institut für Mathematik, Humboldt Universität zu Berlin, e-mail: evgeny@math.hu-berlin.de
[†] Mathematical Faculty, Mordovian State University, 430000 Saransk, e-mail: lobanov@math.mrsu.ru



Figure 1: Zigzag graph Γ^N for (a) $N=3$; (b) $N=1$. The fundamental domain Γ_0 is marked by a bold line.

$x = r_\omega + te_\omega$ and the local coordinate $t \in [0,1]$. We define an oriented zigzag graph Γ^N by identifying $\Gamma_{n,j,k}$ with $\Gamma_{n,j,k+N}$ for all $(n,j,k) \in \mathbb{Z} \times \mathbb{J} \times \mathbb{Z}$. Here and below for simplicity we write $\Gamma_{n,j,k} = \Gamma_{(n,j,k)}$ and $f_{n,j,k} = f_{(n,j,k)}$ etc. Thus Γ^N is a topological space in the quotient topology. As example, the graphs Γ^3 and Γ^1 are illustrated in Fig. 1. We represent Γ^N by

$$\Gamma^N = \cup_{\omega \in \mathcal{Z}} \Gamma_\omega, \qquad \omega = (n,j,k) \in \mathcal{Z} = \mathbb{Z} \times \mathbb{J} \times \mathbb{Z}_N, \quad \mathbb{Z}_N = \mathbb{Z}/(N\mathbb{Z}), \quad \mathbb{J} = \{0,1,2\}.$$

In our paper we assume that $N = 2m+1$ for some integer $m \geqslant 0$. For each function y on Γ^N we define a function $y_\omega = y|_{\Gamma_\omega}, \omega \in \mathcal{Z}$. We identify each function y_ω on Γ_ω with a function on $[0,1]$ by using the local coordinate $t \in [0,1]$ and define the Hilbert space $L^2(\Gamma^N) = \oplus_{\omega \in \mathcal{Z}} L^2(\Gamma_\omega)$. Introduce the space $C(\Gamma^N)$ of continuous functions on Γ^N and the Sobolev space $W^2(\Gamma^N) = \{y, \oplus_{\omega \in \mathcal{Z}} y''_\omega \in L^2(\Gamma^N)$; *$y$ satisfies* **the Kirchhoff Boundary Conditions:** $y \in C(\Gamma^N)$ *and the following identities hold*

$$-y'_{n,0,k}(1) + y'_{n,1,k}(0) - y'_{n,2,k-1}(1) = 0, \qquad y'_{n+1,0,k}(0) - y'_{n,1,k}(1) + y'_{n,2,k}(0) = 0. \tag{1.1}$$

for all $(n,k) \in \mathbb{Z} \times \mathbb{Z}_N$.$\}$ Condition (1.1) means that the sum of derivatives of y at each vertex of Γ^N equals 0 and the orientation of edges gives the sign ±. Our operator H on Γ^N acts in the Hilbert space $L^2(\Gamma^N)$. We define our operator H by $(Hy)_\omega = -y''_\omega + qy_\omega$, where $y = (y_\omega)_{\omega \in \mathcal{Z}} \in \mathfrak{D}(H) = W^2(\Gamma^N)$, $q \in L^2(0,1)$. In the case $q = 0$, we denote the operator H by H^0. The operator H^0 is self-adjoint, see [Ca1], and so is H, see Sect.3.

Recall the needed properties of the Hill operator $\widetilde{H}y = -y'' + q(x)y$ on the real line with a periodic potential $q(x+1) = q(x), x \in \mathbb{R}$. The spectrum of $\widetilde{H}$ is purely absolutely continuous and consists of intervals $\widetilde{\sigma}_n = [\lambda^+_{n-1}, \lambda^-_n], n \geqslant 1$. These intervals are separated



Figure 2: The unrolled honeycomb lattice. The zigzag graph is marked by bold, N=3. The fundamental domain is marked by a double bold line. Each edge Γ_ω has orientation given by $e_\omega = e_j, j = 0, 1, 2$ with the starting point $r_\omega \in \mathbb{R}^2$.

by the gaps $\gamma_n = (\lambda_n^-, \lambda_n^+)$ of length $|\gamma_n| \geqslant 0$. If a gap γ_n is degenerate, i.e. $|\gamma_n| = 0$, then the corresponding segments $\widetilde{\sigma}_n, \widetilde{\sigma}_{n+1}$ merge. For the equation $-y'' + q(x)y = \lambda y$ on the real line we define the fundamental solutions $\vartheta(x, \lambda)$ and $\varphi(x, \lambda), x \in \mathbb{R}$ satisfying $\vartheta(0, \lambda) = \varphi'(0, \lambda) = 1, \vartheta'(0, \lambda) = \varphi(0, \lambda) = 0$. The corresponding monodromy matrix $\mathcal{M}$ and the Lyapunov function Δ are given by

$$\mathcal{M}(\lambda) = \begin{pmatrix} \vartheta(1, \lambda) & \varphi(1, \lambda) \\ \vartheta'(1, \lambda) & \varphi'(1, \lambda) \end{pmatrix}, \qquad \Delta(\lambda) = \frac{\varphi'(1, \lambda) + \vartheta(1, \lambda)}{2}, \quad \lambda \in \mathbb{C}. \tag{1.2}$$

The sequence $\lambda_0^+ < \lambda_1^- \leqslant \lambda_1^+ < .....$ is the spectrum of the equation $-y'' + qy = \lambda y$ with periodic boundary conditions of period 2, that is $y(x + 2) = y(x), x \in \mathbb{R}$. Here equality $\lambda_n^- = \lambda_n^+$ means that $\lambda_n^\pm$ is an eigenvalue of multiplicity 2. Note that $\Delta(\lambda_n^\pm) = (-1)^n$, $n \geqslant 1$. The lowest eigenvalue λ_0^+ is simple, $\Delta(\lambda_0^+) = 1$, and the corresponding eigenfunction has period 1. The eigenfunctions corresponding to $\lambda_n^\pm$ have period 1 if n is even, and they are anti-periodic, that is $y(x + 1) = -y(x)$, $x \in \mathbb{R}$, if n is odd. The derivative of the Lyapunov function has a zero λ_n in each "closed gap" $[\lambda_n^-, \lambda_n^+]$, that is $\Delta'(\lambda_n) = 0$. Let $\mu_n, n \geqslant 1$, be the spectrum of the problem $-y'' + qy = \lambda y, y(0) = y(1) = 0$ (the Dirichlet spectrum), and let $\nu_n, n \geqslant 0$, be the spectrum of the problem $-y'' + qy = \lambda y, y'(0) = y'(1) = 0$ (the Neumann spectrum). It is well-known that $\mu_n, \nu_n \in [\lambda_n^-, \lambda_n^+]$ and $\nu_0 \leqslant \lambda_0^+$. Moreover, a potential q is even, i.e., $q \in L^2_{even}(0, 1) = \Big\{ q \in L^2(0, 1) : q(1 - x) = q(x), x \in [0, 1] \Big\}$ iff $|\gamma_n| = |\mu_n - \nu_n|$ for all $n \geqslant 1$. Define the set $\sigma_D = \{\mu_n, n \geqslant 1\}$ and note that $\sigma_D = \{\lambda \in \mathbb{C} : \varphi(1, \lambda) = 0\}$.

For simplicity we shall denote $\Gamma_{\alpha,1} \subset \Gamma^1$ by Γ_α, for $\alpha = (n,j) \in \mathcal{Z}_1 = \mathbb{Z} \times \mathbb{J}$. Thus $\Gamma^1 = \cup_{\alpha\in\mathcal{Z}_1}\Gamma_\alpha$, see Fig 1. In Theorem 1.1 we will show that H is unitarily equivalent to $\oplus_1^N H_k$, where the operator H_k acts in the Hilbert space $L^2(\Gamma^1)$ and is given by

$$(H_k f)_\alpha = -f_\alpha'' + q f_\alpha, \qquad \oplus_{\alpha\in\mathcal{Z}_1} f_\alpha'' \in L^2(\Gamma^1), \tag{1.3}$$

and the vector function $f = (f_\alpha)_{\alpha\in\mathcal{Z}_1}$ satisfies the Kirchhoff conditions:

$$f_{n,0}(1) = f_{n,1}(0) = s^k f_{n,2}(1), \quad f_{n+1,0}(0) = f_{n,1}(1) = f_{n,2}(0), \quad s = e^{i\frac{2\pi}{N}}, \tag{1.4}$$

$$-f_{n,0}'(1) + f_{n,1}'(0) - s^k f_{n,2}'(1) = 0, \quad f_{n+1,0}'(0) - f_{n,1}'(1) + f_{n,2}'(0) = 0. \tag{1.5}$$

We reduce the spectral problem on the graph Γ^1 to some matrix problem on $\mathbb{R}$. In order to describe this we define the fundamental subgraph Γ_0^1 of Γ^1 by $\Gamma_0^1 = \cup_{j=0}^2 \Gamma_{0,j}$, see Fig. 1. On Γ^1, the group $\mathbb{Z}$ acts via $p \circ \Gamma_{n,j}^1 = \Gamma_{n+p,j}^1$, $(n,j,p) \in \mathcal{Z}_1 \times \mathbb{Z}$. Thus Γ_0^1 is a fundamental domain associated with this group action of $\mathbb{Z}$.

For the operator H_k we construct the fundamental solutions $\Theta_k(x,\lambda) = (\Theta_{k,\alpha}(x,\lambda))_{\alpha\in\mathcal{Z}_1}$, and $\Phi_k(x,\lambda) = (\Phi_{k,\alpha}(x,\lambda))_{\alpha\in\mathcal{Z}_1}$, $(x,\lambda) \in \mathbb{R}\times\mathbb{C}$, which satisfy

$$-f_\alpha'' + q f_\alpha = \lambda f_\alpha, \quad \text{the Kirchhoff Boundary Conditions (1.4),(1.5)}, \tag{1.6}$$

$$\Theta_{k,\beta}(0,\lambda) = \Phi_{k,\beta}'(0,\lambda) = 1, \quad \Theta_{k,\beta}'(0,\lambda) = \Phi_{k,\beta}(0,\lambda) = 0, \qquad \beta = (0,0). \tag{1.7}$$

We introduce the monodromy matrix

$$\mathcal{M}_k(\lambda) = \begin{pmatrix} \Theta_{k,\alpha}(0,\lambda) & \Phi_{k,\alpha}(0,\lambda) \\ \Theta_{k,\alpha}'(0,\lambda) & \Phi_{k,\alpha}'(0,\lambda) \end{pmatrix}, \qquad \alpha = (1,0), \tag{1.8}$$

which is determined by Θ_k, Φ_k on the fundamental domain Γ_0^1.

There are two methods to study periodic differential operators. The direct integral analysis usually used for partial differential operators [ReS] gives general information about the spectrum, but no detailed results. The method of ordinary differential operators, based on the Floquet matrix analysis, gives detailed results. Note that there are a lot of open problems [BBK] even for the Schrödinger operator with periodic 2×2 matrix potentials on the real line .

We introduce the monodromy matrix, similar to the case of Schrödinger operators with periodic matrix potentials on the real line, see [YS]. After this, using the approach from [BK],[BBK],[CK] we introduce the Laypunov functions and study the properties of this functions, similar to the case of the Schrödinger operator with a periodic matrix potential on the real line. This is a crucial point of our analysis. Here we essentially use the results and techniques from the papers [BK], [BBK],[CK]. The recent papers [BBK],[CK] are devoted to Schrödinger operators with periodic matrix potentials (a standard case) on the real line. Remark that Carlson [Ca] studied the monodromy operator to analyze the Schrödinger operator on a product of graphs. His results do not cover our case.

We formulate our first preliminary results.

Theorem 1.1. *(i) The operator H is unitarily equivalent to $\oplus_1^N H_k$, where the operator H_k is given by (1.3)-(1.5).*
(ii) For any $\lambda \in \mathbb{C} \setminus \sigma_D$ and $k \in \mathbb{Z}_N$ there exist unique fundamental solutions Θ_k, Φ_k of the system (1.6),(1.7). Moreover, each of the functions $\Theta_k(x, \lambda), \Phi_k(x, \lambda), x \in \Gamma^1$ is meromorphic in $\lambda \in \mathbb{C} \setminus \sigma_D$ and each matrix $\mathcal{M}_k(\lambda)$ satisfies

$$\mathcal{M}_k = \mathcal{R}^{-1}\mathcal{T}_k\mathcal{R}\mathcal{M}, \quad \mathcal{T}_k = \frac{s^{-\frac{k}{2}}}{2c_k}\begin{pmatrix} 2\Delta & 1 \\ 4\Delta^2 - 4c_k^2 & 2\Delta \end{pmatrix}, \quad \mathcal{R} = \begin{pmatrix} 1 & 0 \\ 0 & \varphi(1, \cdot) \end{pmatrix}. \tag{1.9}$$

In particular,

$$\det \mathcal{M}_k = s^{-k}, \quad \operatorname{Tr} \mathcal{M}_k = \frac{2(\Delta_0 + s_k^2)}{s^{\frac{k}{2}} c_k}, \quad \Delta_0 \equiv \frac{\operatorname{Tr} \mathcal{M}_0}{2} = 2\Delta^2 + \frac{\vartheta'(1, \cdot)\varphi(1, \cdot)}{4} - 1, \tag{1.10}$$

where $c_k = \cos \frac{\pi k}{N}$, $s_k = \sin \frac{\pi k}{N} s = e^{i\frac{2\pi}{N}}$. Furthermore, $\mathcal{R}\mathcal{M}_k\mathcal{R}^{-1} = \mathcal{T}_k\mathcal{R}\mathcal{M}\mathcal{R}^{-1}$ is an entire matrix-valued function. In particular, the function $D_k(\tau, \lambda) = \det(\mathcal{M}_k(\lambda) - \tau I_2)$, where I_2 is the 2×2 identity matrix, is entire with respect to $\lambda, \tau \in \mathbb{C}$.

Remark that in contrast to the Schrödinger operator with periodic matrix potential on the real line (see [YS] or [CK]), the monodromy matrix $\mathcal{M}_k$ has poles at the points $\lambda \in \sigma_D$, which are eigenvalues of H_k, see Theorem 1.2. Such a phenomenon has already been observed e.g. in [SA], [MV], [Ku].

Define the subspace $\mathcal{H}_k(\lambda) = \{\psi \in \mathfrak{D}(H_k) : H_k\psi = \lambda\psi\}$ for $\lambda \in \sigma_{pp}(H_k), k \in \mathbb{Z}_N$. If $\dim \mathcal{H}_k(\lambda_0) = \infty$ for some $\lambda_0 \in \sigma_{pp}(H_k)$, then we say that $\{\lambda_0\}$ is a flat band. In Theorem 1.2 we describe all flat bands and supports of eigenfunctions (see Fig. 3).

Theorem 1.2. *Let $(\lambda, k) \in \sigma_D \times \mathbb{Z}_N$. Then*
(i) Each eigenfunction from $\mathcal{H}_k(\lambda)$ vanishes at all vertices of Γ^1.
(ii) Let $\varphi = \varphi(\cdot, \lambda)|_{[0,1]}$ and put $c = \varphi'(1, \lambda)$. Define the function $\psi^{(0)} = (\psi_\alpha^{(0)})_{\alpha \in \mathcal{Z}_1}$ by :
if $\eta = 1 - s^k c^2 \neq 0$, then

$$\psi_{n,j}^{(0)} = 0, \text{for all } n \neq 0, -1, \ j \in \mathbb{Z}_3, \qquad \text{and} \quad \psi_{0,0}^{(0)} = \eta\varphi, \ \psi_{0,1}^{(0)} = c\varphi, \ \psi_{0,2}^{(0)} = c^2\varphi,$$
$$\psi_{-1,0}^{(0)} = 0, \ \psi_{-1,1}^{(0)} = -s^k c\varphi, \ \psi_{-1,2}^{(0)} = -\varphi, \tag{1.11}$$

if $\eta = 0$, then

$$\psi_{0,0}^{(0)} = 0, \quad \psi_{0,1}^{(0)} = \varphi, \quad \psi_{0,2}^{(0)} = c\varphi, \quad \psi_{n,j}^{(0)} = 0, \ \text{all } n \neq 0, j \in \mathbb{Z}_3. \tag{1.12}$$

Then each $\psi^{(n)} = (\psi_{n-m,j}^{(0)})_{(m,j) \in \mathcal{Z}_1} \in \mathcal{H}_k(\lambda), n \in \mathbb{Z}$ and each $f \in \mathcal{H}_k(\lambda)$ has the form

$$f = \sum_{n \in \mathbb{Z}} \widehat{f}_n \psi^{(n)}, \qquad \widehat{f}_n = \begin{cases} \eta^{-1} f'_{n,0}(0) & \text{if } \eta \neq 0 \\ f'_{n,1}(0) & \text{if } \eta = 0 \end{cases}, \qquad (\widehat{f}_n)_{n \in \mathbb{Z}} \in \ell^2. \tag{1.13}$$

Moreover, the mapping $f \to \{\widehat{f}_n\}_{n \in \mathbb{Z}}$ is a linear isomorphism between $\mathcal{H}_k(\lambda)$ and ℓ^2.



(a) (b)

Figure 3: The supports of the eigenfunction $\psi^{(0)}$: (a) $\eta = 1 - s^k\varphi'(1,\lambda)^2 = 0$; (b) $\eta \neq 0$.

Let $\tau_{k,\pm}$ be the eigenvalues of $\mathcal{M}_k, k \in \mathbb{Z}_N$. Using (1.10) we deduce that

$$\tau_{k,-}\tau_{k,+} = s^{-k}, \quad \tau_{k,-} + \tau_{k,+} = \operatorname{Tr}\mathcal{M}_k = s^{-k}\operatorname{Tr}\mathcal{M}_{-k}, \quad \tau_{-k,+} = \tau_{k,-}s^k, \quad \tau_{-k,-} = \tau_{k,+}s^k. \tag{1.14}$$

If $k = 0$, then we introduce the standard Lyapunov function $\Delta_0 = \frac{\operatorname{Tr}\mathcal{M}_0}{2} = \frac{1}{2}(\tau_{0,+} + \frac{1}{\tau_{0,+}})$, which is entire. If $k \neq 0$, then we define the Lyapunov functions $\Delta_{k,\pm} = \frac{1}{2}(\tau_{k,\pm} + \frac{1}{\tau_{k,\pm}})$ (see [BBK],[CK]) and using (1.14) we get $\Delta_{-k,\pm} = \Delta_{k,\pm}$. Below we prove the following identities

$$\Delta_{k,\pm} = \xi_k \pm \sqrt{\rho_k}, \qquad \xi_k = \Delta_0 + s_k^2, \quad \rho_k = \frac{s_k^2}{c_k^2}(c_k^2 - \xi_k^2), \quad c_k = \cos\frac{\pi k}{N}, \; s_k = \sin\frac{\pi k}{N}. \tag{1.15}$$

If $q = 0$, then we denote the corresponding functions by $\Delta_k^0, \rho_k^0, ...$ In particular, we have

$$\Delta_0^0 = \frac{9\cos 2\sqrt{\lambda} - 1}{8}, \qquad \xi_k^0 = \Delta_0^0 + s_k^2, \quad \rho_k^0 = \frac{s_k^2}{c_k^2}(c_k^2 - (\xi_k^0)^2), \tag{1.16}$$

Introduce the two sheeted Riemann surface $\mathcal{R}_k$ (of infinite genus) defined by $\sqrt{\rho_k}$. The functions $\Delta_{k,\pm}, k \in \overline{m} = \{1,2,..,m\}$, are the branches of $\Delta_k = \xi_k + \sqrt{\rho_k}$ (see Fig.4) on the Riemann surface $\mathcal{R}_k$. We describe spectral properties of H_k in terms of Δ_k.

Theorem 1.3. *(i) The Lyapunov functions $\Delta_{k,\pm}, k \in \overline{m}$ satisfy (1.15).*
(ii) For any $k \in \mathbb{Z}_N$ the following identities hold:

$$\sigma(H_k) = \sigma_{pp}(H_k) \cup \sigma_{ac}(H_k), \quad \sigma_{pp}(H_k) = \sigma_D, \quad \sigma_{ac}(H_k) = \{\lambda \in \mathbb{R} : \Delta_k(\lambda) \in [-1,1]\}. \tag{1.17}$$

(iii) If $\Delta_k(\lambda) \in (-1,1)$ for some $\lambda \in \mathbb{R}, k = 0,..,m$ and λ is not a branch point of $\Delta_k(\lambda)$, then $\Delta_k'(\lambda) \neq 0$.

Remark. 1) If we know Δ_0, then we determine all $\Delta_k, \rho_k, k \in \mathbb{Z}_N$ by (1.15). 2) If we know ρ_j for some $j \in \overline{m}$, then by (1.15), we determine all $\Delta_k, \rho_k, k \in \mathbb{Z}_N$.

Definition 1. *A zero of $\rho_k, k \in \overline{m} = \{1,2,..,m\}$ is called a* **resonance** *of H.*

Let $\lambda_{0,2n}^{\pm}$ and $\lambda_{0,2n+1}^{\pm}, n \geqslant 0$ be the zeros of $\det(\mathcal{M}_0 - I_2)$ and $\det(\mathcal{M}_0 + I_2)$ respectively (counted with multiplicity). In Theorem 1.4 we will show that the periodic eigenvalues $\lambda_{0,2n}^{\pm}$ and the anti-periodic eigenvalues $\lambda_{0,2n+1}^{\pm}$ satisfy

$$\Delta_0(\lambda_{0,n}^{\pm}) = (-1)^n, \quad \lambda_{0,0}^+ < \lambda_{0,1}^- < \lambda_{0,1}^+ < \lambda_{0,2}^- \leqslant \lambda_{0,2}^+ < \lambda_{0,3}^- < \lambda_{0,3}^+ < \lambda_{0,4}^- \leqslant \lambda_{0,4}^+ < .... \tag{1.18}$$



Figure 4: Typical Lyapunov functions and the zigzag nanotube. The function Δ_3 is shown for $q \in L_{\text{even}}(0,1), N = 6$.

For H_0 we introduce the spectral bands $\sigma_{0,n} = [\lambda^+_{0,n-1}, \lambda^-_{0,n}]$ and the gaps $\gamma_{0,n} = (\lambda^-_{0,n}, \lambda^+_{0,n})$, $n \geqslant 1$. Using the asymptotics of the fundamental solutions ϑ, φ, we obtain

$$\Delta_0(\lambda) = \Delta^0_0(\lambda) + \frac{9q_0}{8}\frac{\sin 2\sqrt{\lambda}}{\sqrt{\lambda}} + \frac{O(e^{|\operatorname{Im}\sqrt{\lambda}|})}{|\lambda|}, \qquad q_0 = \int_0^1 q(t)dt \tag{1.19}$$

as $|\lambda| \to \infty$. The following theorem describes the basic properties of H_0.

Theorem 1.4. *The function Δ_0 is entire and has the following properties:*
(i) The function Δ'_0 has only real simple zeros $\lambda_{0,n}, n \geqslant 1$, which are separated by the simple zeros $\eta_{0,n}$ of Δ_0: $\eta_{0,1} < \lambda_{0,1} < \eta_{0,2} < \lambda_{0,2} < \eta_{0,3} < ...$ and satisfy

$$\nu_n, \mu_n \in \gamma_{0,2n}, \quad \Delta_0(\lambda_{0,2n}) \geqslant 1, \qquad \Delta_0(\lambda_{0,2n-1}) \leqslant -\frac{5}{4}, \qquad \text{for any } n \geqslant 1. \tag{1.20}$$

(ii) The periodic and anti-periodic eigenvalues $\lambda^\pm_{0,n}, n \geqslant 0$ satisfy (1.18) and have asymptotics

$$\lambda^\pm_{0,1+2n} = (\pi\frac{2n+1}{2} \pm \phi)^2 + q_0 + o(n^{-1}), \qquad \lambda^\pm_{2n} = (\pi n)^2 + q_0 \pm \left||\hat{q}_n|^2 - \frac{\hat{q}^2_{sn}}{9}\right|^{\frac{1}{2}} + \frac{O(1)}{n}, \tag{1.21}$$

where $\phi = \arcsin\frac{1}{3} \in [0, \frac{\pi}{2}]$ and $\hat{q}_n = \int_0^1 q(t)e^{i2\pi nt}dt, \hat{q}_{sn} = \operatorname{Im}\hat{q}_n$. Moreover, $(-1)^n\Delta_0(\lambda) \geqslant 1$ for all $\lambda \in [\lambda^-_{0,n}, \lambda^+_{0,n}]$ and $\lambda_{0,n} \in [\lambda^-_{0,n}, \lambda^+_{0,n}]$.

(iii) $|\gamma_n| = |\mu_n - \nu_n|$ *iff* $\gamma_n = \gamma_{0,2n}$.
(iv) $\gamma_{0,2n} \subset \gamma_n$ *for all* $n \geqslant 1$. *Moreover, for fixed* $n \geqslant 1$ *we have* $\gamma_{0,2n} = \emptyset$ *iff* $\gamma_n = \emptyset$.
(v) $\Delta_0(\lambda_{1+2n}) = -\frac{5}{4}$ *for all* $n \geqslant 0$ *iff* $q \in L^2_{even}(0,1)$.

Remark that the last theorem gives the asymptotics of the gap length.

Let $\lambda^{\pm}_{k,2n}$ and $\lambda^{\pm}_{k,2n+1}, n \geqslant 0$ be the zeros of $\det(\mathcal{M}_k - I_2)$ and $\det(\mathcal{M}_k + I_2)$. Below we will show that the periodic eigenvalues $\lambda^{\pm}_{k,2n}$ and the anti-periodic eigenvalues $\lambda^{\pm}_{k,2n+1}$ satisfy the equations $\Delta_k(\lambda^{\pm}_{k,n}) = (-1)^n$. In Theorem 1.5 we show that $\lambda^{\pm}_{k,n}$ satisfy the equation

$$\Delta_0(\lambda^{\pm}_{k,2n}) = \cos\frac{2\pi k}{N}, \qquad \Delta_0(\lambda^{\pm}_{k,2n+1}) = -1, \qquad k \in \overline{m}, \tag{1.22}$$

and the labeling is given by: each $\lambda^{\pm}_{k,n}$ is double and

$$\lambda^+_{0,0} < \lambda^+_{1,0} < \lambda^+_{2,0} < ... < \lambda^+_{m,0} < \lambda^-_{0,1} < \lambda^+_{0,1} < \lambda^-_{m,2} < \lambda^-_{m-1,2} < ... < \lambda^-_{0,2} \leqslant \lambda^+_{0,2} < ..., \tag{1.23}$$

$$\lambda^{\pm}_{0,2n-1} = \lambda^{\pm}_{k,2n-1}, \quad (k,n) \in \overline{m} \times \mathbb{N}. \tag{1.24}$$

The periodic eigenvalues $\lambda^{\pm}_{k,2n}, k \in \overline{m} = \{1,..,m\}, n \geqslant 0$ (i.e., $\Delta_k(\lambda^{\pm}_{k,2n}) = 1$) satisfy

$$\lambda^{\pm}_{k,2n} = (\pi n \pm \phi_k)^2 + q_0 + \frac{o(1)}{n}, \qquad \phi_k = \frac{1}{2}\arccos\frac{1+8c_{2k}}{9} \in [0, \frac{\pi}{2}] \quad as \quad n \to \infty. \tag{1.25}$$

Let $r^{\pm}_{k,n}, k \in \overline{m}, n \geqslant 0$ be the zeros of ρ_k. Below we will show that $r^{\pm}_{k,n}$ satisfy the equation

$$\Delta_0(r^{\pm}_{k,2n}) = c_k - s_k^2, \qquad \Delta_0(r^{\pm}_{k,2n+1}) = -c_k - s_k^2, \qquad k \in \overline{m}, \tag{1.26}$$

they are real and the labeling is given by

$$r^+_{k,0} < r^-_{k,1} < r^+_{k,1} < r^-_{k,2} < r^+_{k,2} < .., \qquad k \neq \frac{N}{3}, \tag{1.27}$$

$$r^+_{k,0} < r^-_{k,1} \leqslant r^+_{k,1} < r^-_{k,2} < r^+_{k,2} < r^-_{k,3} \leqslant r^+_{k,3}.., \qquad k = \frac{N}{3}. \tag{1.28}$$

The resonances $r^{\pm}_{k,2n+s}, s = 0,1$ (for $k \in \overline{m}, k \neq \frac{N}{3}$) satisfy

$$r^{\pm}_{k,n} = (\frac{\pi n}{2} \pm b_{k,n})^2 + q_0 + \frac{o(1)}{n}, \qquad b_{k,2n} = \phi_{k,0}, \quad b_{k,2n+1} = \frac{\pi}{2} - \phi_{k,1}, \tag{1.29}$$

$\phi_{k,s} = \frac{1}{2}\arccos\frac{1+(-1)^s 8c_k - 8s_k^2}{9} \in [0, \frac{\pi}{2}]$ as $n \to \infty$. We describe the spectral properties of H.

Theorem 1.5. *(i) The periodic and anti-periodic eigenvalues* $\lambda^{\pm}_{k,n}, k \in \overline{m} = \{1,..,m\}, n \geqslant 0$ *satisfy Eq. (1.22) and the relations (1.23)- (1.25).*
(ii) The resonances $r^{\pm}_{k,n}, k \in \overline{m}, n \geqslant 0$ *satisfy Eq. (1.26) and estimates (1.27)-(1.29).*
(iii) For $n \geqslant 1, k \in \overline{m}$ *the following identities are fulfilled:*

$$\sigma_{ac}(H) = \cup_{n\geqslant 1} S_n = \cup_0^N \sigma_{ac}(H_k), \quad S_n = [E^+_{n-1}, E^-_n] = \cup_{k=0}^m \sigma_{k,n}, \quad \cap_{k=0}^m \sigma_{k,n} \neq \emptyset, \tag{1.30}$$

$$\sigma_{0,n}=[\lambda_{0,n-1}^+,\lambda_{0,n}^-],\quad \text{and}\quad \lambda_{k,n}^{\pm}\in\sigma_{k,n}=[r_{k,n-1}^+ r_{k,n}^-],\tag{1.31}$$

$$G_n=(E_n^-,E_n^+)=\cap_0^m\gamma_{k,n},\quad G_{2n}=\gamma_{0,2n}.\tag{1.32}$$

(iv) If $n\geqslant 1$ *is fixed, then* $G_{2n}=\emptyset$ *iff* $\gamma_n=\emptyset$. *Moreover,* $|G_{2n}|=o(1)$ *as* $n\to\infty$.
(v) If $\frac{N}{3}\in\mathbb{Z}$, *then* $G_{2n+1}=\emptyset$ *for all* $n\geqslant 0$ *iff* $q\in L_{even}(0,1)$. *Moreover, each odd gap* G_n *has the form* $G_n=(r_{p,n}^-,r_{p,n}^+)$ *(*$n\geqslant 1$ *is odd) for some integer* $p=p(n)$ *and* $r_{p,n}^{\pm}$ *satisfy*

$$r_{p,n}^{\pm}=\pi^2\frac{n^2}{4}+q_0\pm|\widetilde{q}_{cn}|+o(n^{-1})\quad as\quad n\to\infty,\quad \widetilde{q}_{cn}=\int_0^1 q(t)\cos\pi n t dt.\tag{1.33}$$

(vi) If $\frac{N}{3}\notin\mathbb{Z}$, *then each gap* $G_{2n+1}\neq\emptyset, n\geqslant 0$ *and* $|G_{2n+1}|\to\infty$ *as* $n\to\infty$.

Note that in the armchair case (see [BBKL]) there exist non-real resonances for some specific potentials. That all resonances are real is a peculiarity of the high symmetry of a zigzag periodic graph.

Finally, we formulate our results about **the inverse problem**. By Theorem 1.3, $E_n=\mu_n, n\geqslant 1$ are the eigenvalues of H and they satisfy (see [PT]) $E_n=\pi^2n^2+q_0+\varkappa_n$, where $q_0=\int_0^1 q(t)dt$ and $(\varkappa_n)_1^\infty\in\ell^2=\ell_0^2$. Here $\pi^2n^2, n\geqslant 1$ are the unperturbed eigenvalues and the real Hilbert spaces ℓ_p^2 is given by $\ell_p^2=\biggl\{f=(f_n)_1^\infty:\sum_{n\geqslant 1}n^{2p}f_n^2<\infty\biggr\}$, $p\geqslant 0$. The monotonicity property $E_1<E_2<...$ gives that if q runs through $L^2(0,1)$, then $(\varkappa_n)_1^\infty$ doesn't run through the whole space ℓ^2. In order to describe this situation, we introduce the open and convex set $\mathcal{K}=\{(\varkappa_n)_1^\infty\in\ell^2:\pi^2+\varkappa_1<(2\pi)^2+\varkappa_2<\ldots\}\subset\ell^2$. Let $Hf_n=E_nf_n$ for some nonzero eigenfunction $f_n\in\mathcal{H}(E_n)$. Then $f_{n,\omega}\neq 0$ for some $\omega\in\mathcal{Z}$ and using Theorem 1.2 we obtain

$$f_{n,\omega}(0)=f_{n,\omega}(1)=0,\quad f_{n,\omega}'(0)=C_{n,\omega},\quad f_{n,\omega}'(1)^2=C_{n,\omega}^2e^{2h_n}>0,\quad h_n\in\mathbb{R},\quad any\quad n\geqslant 1,$$

for some constant $C_{n,\omega}\neq 0$. From Theorem 1.3-1.4 we have a simple corollary.

Corollary 1.6. *(i) The operator* H *has only a finite number of non degenerate gaps* G_n *iff* $\frac{N}{3}\in\mathbb{Z}$ *and* q *is an even finite gap potential for the operator* $-y''+qy$ *on the real line.*
(ii) The mapping $\Phi:q\to\biggl(q_0,(\varkappa_n)_1^\infty;(h_n)_1^\infty\biggr)$ *is a real-analytic isomorphism between* $L^2(0,1)$ *and* $\mathbb{R}\times\mathcal{K}\times\ell_1^2$, *where the set* $\mathcal{K}=\{(\varkappa_n)_1^\infty\in\ell^2:\pi^2+\varkappa_1<(2\pi)^2+\varkappa_2<\ldots\}\subset\ell^2$.
(iii) The mapping $\Phi_e:q\to\biggl(q_0,(\varkappa_n)_1^\infty\biggr)$ *is a real-analytic isomorphism between* $L_{even}^2(0,1)$ *and* $\mathbb{R}\times\mathcal{K}$.

For the convenience of the reader we briefly describe the structure of carbon nanotubes (see [Ha], [SDD]) and explain how they are related to the graph Γ^N. Graphene is a single 2D layer of graphite forming a honeycomb lattice, as in Fig. 1 and 2. A carbon nanotube is a honeycomb lattice "rolled up" into a cylinder. In carbon nanotubes, the graphene sheet



Figure 5: The unrolled honeycomb lattice of nanotube. The unit cell is spanned by the vectors Ω_1 and Ω_2. The type of the nanotube is defined by the pair $(N_1, N_2) \in \mathbb{N}^2$, $N_1 \geqslant N_2$, and corresponding chiral vector $\Omega = N_1\Omega_1 + N_2\Omega_2$.

is "rolled up" in such a way that a so-called chiral vector $\Omega = N_1\Omega_1 + N_2\Omega_2$ becomes the circumference of the tube, where $\Omega_1 = e_1 + e_2, \Omega_2 = e_0 + e_1$ see Fig 5 and 2. The chiral vector Ω, which is usually denoted by the pair of integers (N_1, N_2), uniquely defines a particular tube. Tubes of type $(N, 0)$ are called zigzag tubes, see Fig 5 and 4. They exhibit a zigzag pattern along the circumference, see Figure 1. (N, N)-tubes are called armchair tubes.

Remark that in this zigzag standard physical model only each vertical edge has length 1 and the other edges are shorter (or might be shorter). In this paper, we avoid this additional technical difficulty. Instead, we do our detailed spectral analysis for the simplest possible model of a physical nanotube T^N (to be described below and see Fig. 4). This will simplify our analysis (in another paper [KL]) of the Schrödinger operator $\mathscr{H}_B = (-i\nabla - \mathscr{A})^2 + q$ on T^N with a periodic potential q and a uniform magnetic field $\mathscr{B} = B(0, 0, 1) \in \mathbb{R}^3$, $B \in \mathbb{R}$. The corresponding vector potential is given by $\mathscr{A}(x) = \frac{1}{2}[\mathscr{B}, x] = \frac{B}{2}(-x_2, x_1, 0)$, $x = (x_1, x_2, x_3) \in \mathbb{R}^3$. Maybe the second simplest model is the armchair nanotube [Ha], but then the monodromy matrix is 4×4 and there exist complex resonances for some specific potential q (see [BBKL]).

Finally we consider the Schrödinger operator $\mathscr{H} = -\partial^2 + q$ with a periodic potential q on the zigzag nanotube $T^N \subset \mathbb{R}^3$. Our model nanotube T^N is a union of edges T_ω of length 1, i.e., $T^N = \cup_{\omega\in\mathcal{Z}} T_\omega$, see Fig. 4. Each edge $T_\omega = \{x = \mathbf{r}_\omega + t\mathbf{e}_\omega,\ t \in [0, 1]\}$ is oriented by the vector $\mathbf{e}_\omega \in \mathbb{R}^3$ and has starting point $\mathbf{r}_\omega \in \mathbb{R}^{\mathbf{3}}$. We have the coordinate $x = \mathbf{r}_\omega + t\mathbf{e}_\omega$ and the local coordinate $t \in [0, 1]$ (length preserving). We define $\mathbf{r}_\omega, \mathbf{e}_\omega, \omega = (n, j, k) \in \mathcal{Z}$ by

$$\mathbf{r}_{n,0,k} = \varkappa_{n+2k} + \frac{3n}{2}\mathbf{e}_0, \quad \mathbf{r}_{n,1,k} = \mathbf{r}_{n,0,k} + \mathbf{e}_0, \quad \mathbf{r}_{n,2,k} = \mathbf{r}_{n+1,0,k},$$

$$\mathbf{e}_{n,0,k} = \mathbf{e}_0, \quad \mathbf{e}_{n,1,k} = \varkappa_{n+2k+1} - \varkappa_{n+2k} + \frac{\mathbf{e}_0}{2}, \quad \mathbf{e}_{n,2,k} = \varkappa_{n+2k+2} - \varkappa_{n+2k+1} - \frac{\mathbf{e}_0}{2},$$

where $\varkappa_j = R_N(c_j, s_j, 0) \in \mathbb{R}^3$, $R_N = \frac{\sqrt{3}}{4\sin\frac{\pi}{2N}}$, The points $\mathbf{r}_{0,0,k}$ are vertices of the regular N-gon $\mathscr{P}_0$. The vertical edge $T_{0,0,k}$ is the segment and lies on the cylinder $\mathcal{C} \equiv \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1^2 + x_2^2 = R_N^2\}$. The starting points $\mathbf{r}_{1,0,k} = \mathbf{r}_{0,2,k} = \varkappa_{1+2k} + \frac{3}{2}\mathbf{e}_0$, $k \in \mathbb{Z}_N$ are the vertices of the regular N-gon $\mathscr{P}_1$. $\mathscr{P}_1$ arises from $\mathscr{P}_0$ by the following motion: rotate around the axis of the cylinder $\mathcal{C}$ by the angle $\frac{\pi}{N}$ and translate by $\frac{3}{2}e_0$. The non-vertical edges $T_{0,1,k}$ and $T_{0,2,k}$ have positive and negative projections on the vector $\mathbf{e}_0$. Repeating this procedure we obtain all edges of T^N. Note that each non-vertical edge $T_{0,j,k}, j = 1, 2$ (without the endpoints) lies inside the cylinder $\mathcal{C}$.

Our operator $\mathscr{H}$ on T^N acts in the space $L^2(T^N) = \oplus_{\omega\in\mathcal{Z}} L^2(T_\omega)$. Then acting on the edge T_ω, $\mathscr{H}$ is the ordinary differential operator given by $(\mathscr{H}f)_\omega = -f_\omega'' + qf_\omega$, $f_\omega, f_\omega'' \in L^2(\Gamma_\omega)$, $\omega \in \mathcal{Z}$ where $q \in L^2(0,1)$ and $f \in \mathfrak{D}(\mathscr{H}) = W^2(T^N)$ satisfies the Kirchhoff Conditions (1.1). Thus the operator $\mathscr{H}$ on T^N is unitarily equivalent to the operator H on Γ^N.

We remark that such models were introduce by Pauling [Pa] in 1936 to simulate aromatic molecules. They were described in more detail by Ruedenberg and Scherr [RS] in 1953. For further physical models see [Ha], [SDD]. For papers on spectral analysis of such operators, see the references in [Ku].

We present the plan of the paper. In Sect. 2 we construct the fundamental solutions and describe the basic properties of the monodromy operator. Moreover, we prove Theorem 1.2 about the eigenfunctions. In Sect. 3 we prove the basic properties of the Lyapunov function Δ_0 which is entire on the plane. This function is important to study the functions $\Delta_k, k \in \mathbb{Z}_N$ which are analytic on the two-sheeted Riemann surface $\mathcal{R}_k$. In Sect. 4 we prove the basic properties of the Lyapunov function $\Delta_k, k \in \mathbb{Z}_N$ and Corollary 1.6.

2 Fundamental solutions and eigenfunctions

Proof of Theorem 1.1. (i) Define the operator $\mathcal{S}$ in $\mathbb{C}^N$ by $\mathcal{S}u = (u_N, u_1, \dots, u_{N-1})^\top$, $u = (u_n)_1^N \in \mathbb{C}^N$. The unitary operator $\mathcal{S}$ has the form $\mathcal{S} = \sum_1^N s^k \mathcal{P}_k$, where $\mathcal{S}v_k = s^k v_k$ and $v_k = \frac{1}{N^{\frac{1}{2}}}(1, s^{-k}, s^{-2k}, ..., s^{-kN+k})$ is an eigenvector (recall $s = e^{i\frac{2\pi}{N}}$); $\mathcal{P}_k u = v_k(u, v_k)$ is a projector. The function f in the Kirchhoff boundary conditions (1.1) is a vector function $f = (f_\omega), \omega = (n, j, k) \in \mathcal{Z}$. We define a new vector-valued function $f_{n,j} = (f_{n,j,k})_k^N$, where each $f_{n,j}, (n, j) \in \mathcal{Z}_1 = \mathbb{Z} \times \{0, 1, 2\}$ is an $\mathbb{C}^N-$ vector, which satisfies the equation $-f_{n,j}'' + qf_{n,j} = \lambda f_{n,j}$, and the conditions (which follow from the Kirchhoff conditions (1.1))

$$f_{n,0}(1) = f_{n,1}(0) = \mathcal{S}f_{n,2}(1), \quad f_{n+1,0}(0) = f_{n,1}(1) = f_{n,2}(0), \tag{2.1}$$

$$-f_{n,0}'(1) + f_{n,1}'(0) - \mathcal{S}f_{n,2}'(1) = 0, \quad f_{n+1,0}'(0) - f_{n,1}'(1) + f_{n,2}'(0) = 0, \tag{2.2}$$

for all $n \in \mathbb{Z}$. The operators $\mathcal{S}$ and H commute, then $H = \oplus_1^N (H\mathcal{P}_k)$. Using (2.1), (2.2) we deduce that $H\mathcal{P}_k$ is unitarily equivalent to the operator H_k. The operator H_k on $\Gamma^1 = \cup_{\alpha\in\mathcal{Z}_1}\Gamma_\alpha^1$ acts in the Hilbert space $L^2(\Gamma^1)$. Then acting on the edge Γ_α, the operator H_k given by: $(H_k f)_\alpha = -f_\alpha'' + q(t)f_\omega$, where $f_\alpha, f_\alpha'' \in L^2(0,1), \alpha = (n, j) \in \mathcal{Z}_1$, on the vector functions $f = (f_\alpha)_{\alpha \in \mathcal{Z}_1}$, which satisfy the boundary conditions (1.4), (1.5), i.e.,

$$f_{n,0}(1) = f_{n,1}(0) = s^k f_{n,2}(1), \quad f_{n+1,0}(0) = f_{n,1}(1) = f_{n,2}(0), \quad s = e^{i\frac{2\pi}{N}}, \tag{2.3}$$

$$-f'_{n,0}(1) + f'_{n,1}(0) - s^k f'_{n,2}(1) = 0, \qquad f'_{n+1,0}(0) - f'_{n,1}(1) + f'_{n,2}(0) = 0. \tag{2.4}$$

(ii) Fix $k \in \mathbb{Z}_N$. We determine the fundamental solutions $\Theta_k(x,\lambda), \Phi_k(x,\lambda)$ of the equation $-f'' + qf = \lambda f, \quad \lambda \in \mathbb{C}$, on Γ^1, where Θ_k, Φ_k satisfy (2.3)-(2.4) and

$$\Theta_{k,\alpha}(0,\lambda) = \Phi'_{k,\alpha}(0,\lambda) = 1, \quad \Theta'_{k,\alpha}(0,\lambda) = \Phi_{k,\alpha}(0,\lambda) = 0, \qquad \alpha = (0,0). \tag{2.5}$$

Below we assume $f = \Theta_k$ or $f = \Phi_k$ and let $\vartheta_t = \vartheta(1,\lambda), \varphi_t = \varphi(t,\lambda), ...$ Any solution of the equation $-f'' + qf = \lambda f$ satisfies $f(t) = \vartheta_t f(0) + \frac{\varphi_t}{\varphi_1}(f(1) - \vartheta_1 f(0)), \quad t \in [0,1]$ and

$$f'(0) = \frac{f(1) - \vartheta_1 f(0)}{\varphi_1}, \quad f'(1) = \frac{\varphi'_1 f(1) - f(0)}{\varphi_1}. \tag{2.6}$$

Substituting (2.6) for $f = f_{01}, f = f_{02}$ into the first Eq. in (2.4) at $n = 0$, we obtain

$$-\varphi_1 f'_{0,0}(1) + (f_{0,1}(1) - \vartheta_1 f_{0,1}(0)) - s^k(\varphi'_1 f_{0,2}(1) - f_{0,2}(0)) = 0.$$

Then the condition (2.3) gives

$$\varphi_1 f'_{0,0}(1) + (f_{1,0}(0) - \vartheta_1 f_{0,0}(1)) - s^k(\varphi'_1 s^{-k} f_{0,0}(1) - f_{1,0}(0)) = 0, \tag{2.7}$$

which implies

$$-\varphi_1 f'_{0,0}(1) + u f_{1,0}(0) - 2\Delta f_{0,0}(1) = 0, \qquad u = 1 + s^k = 2c_k s^{\frac{k}{2}}. \tag{2.8}$$

Thus we obtain **the first basic identity**

$$f_{1,0}(0) = \frac{\varphi_1}{u} f'_{0,0}(1) + \frac{2\Delta}{u} f_{0,0}(1), \qquad u = 2c_k s^{\frac{k}{2}}, \quad c_k = \cos\frac{\pi k}{N}. \tag{2.9}$$

Substituting (2.6) for $f = f_{01}, f = f_{02}$ into the second Eq. in (2.4) at $n = 0$, we get

$$\varphi_1 f'_{1,0}(0) - (\varphi'_1 f_{0,1}(1) - f_{0,1}(0)) + (f_{0,2}(1) - \vartheta_1 f_{0,2}(0)) = 0.$$

Using (2.3), we obtain

$$\varphi_1 f'_{1,0}(0) - (\varphi'_1 f_{1,0}(0) - f_{0,0}(1)) + (s^{-k} f_{0,0}(1) - \vartheta_1 f_{1,0}(0)) = 0, \tag{2.10}$$

which yields

$$\varphi_1 f'_{1,0}(0) = 2\Delta f_{1,0}(0) - u_2 f_{0,0}(1), \qquad u_2 = 1 + s^{-k} = 2c_k s^{-\frac{k}{2}}. \tag{2.11}$$

Then substituting (2.9) into (2.11), we obtain **the second basic identity**

$$f_{1,0}'(0) = \frac{4(\Delta^2 - c_k^2)}{u\varphi_1} f_{0,0}(1) + \frac{2\Delta}{u} f_{0,0}'(1), \qquad u = 2c_k s^{\frac{k}{2}}. \tag{2.12}$$

Substituting consequently $f = \Theta_k$ and $f = \Phi_k$ into (2.9) and into (2.12), and after this $\Theta_{k,\omega}(0,\lambda), \Theta_{k,\omega}'(0,\lambda), ..$ into $\mathcal{M}_k(\lambda) = \begin{pmatrix} \Theta_{k,\alpha}(0,\lambda) & \Phi_{k,\alpha}(0,\lambda) \\ \Theta_{k,\alpha}'(0,\lambda) & \Phi_{k,\alpha}'(0,\lambda) \end{pmatrix}$, $\alpha = (1,0)$, we obtain (1.9).

Using (1.9) we obtain $\det \mathcal{T}_k = \frac{s^{-k} 4c_k^2}{4c_k^2} = s^{-k}$, $\det \mathcal{M}_k = \det \mathcal{T}_k \mathcal{M} = s^{-k}$, since $\det \mathcal{M} = 1$. Moreover, using $\mathcal{R}\mathcal{M}\mathcal{R}^{-1} = \begin{pmatrix} \vartheta_1 & 1 \\ \varphi_1 \vartheta_1' & \varphi_1' \end{pmatrix}$, $\quad \mathcal{R} = \begin{pmatrix} 1 & 0 \\ 0 & \varphi_1 \end{pmatrix}$, we obtain

$$\operatorname{Tr} \mathcal{M}_k = \operatorname{Tr} \mathcal{T}_k \mathcal{R}\mathcal{M}\mathcal{R}^{-1} = \frac{2s^{-\frac{k}{2}}}{c_k} \operatorname{Tr} \begin{pmatrix} 2\Delta & 1 \\ 4\Delta^2 - 4c_k^2 & 2\Delta \end{pmatrix} \begin{pmatrix} \vartheta_1 & 1 \\ \varphi_1 \vartheta_1' & \varphi_1' \end{pmatrix}$$

$$= \frac{2s^{-\frac{k}{2}}}{c_k} \Big(2\Delta\vartheta_1 + \varphi_1\vartheta_1' + 4(\Delta^2 - c_k^2) + 2\Delta\varphi_1' \Big) = \frac{2s^{-\frac{k}{2}}}{c_k} \Big(2\Delta^2 + \frac{\varphi_1\vartheta_1'}{4} - c_k^2 \Big) = \frac{2s^{-\frac{k}{2}}(\Delta_0 + s_k^2)}{c_k}$$

which gives (1.10). ■

Recall that $H^0, \mathfrak{D}(H^0) = W^2(\Gamma^N)$ is self-adjoint [Ca1]. Let $\|f\|^2 = \int_{\Gamma^N} |f(x)|^2 dx$ and let $\|f_\omega\|_0^2 = \int_0^1 |f_\omega(t)|^2 dt$ for $f_\omega \in L^2(\Gamma_\omega), \omega \in \mathcal{Z}$. Assume that

$$\|qf\|^2 \leqslant \frac{1}{2}\|H^0 f\|^2 + C\|f\|^2, \quad all \quad f \in \mathfrak{D}(H^0), \tag{2.13}$$

for some constant $C > 0$. Then by the Kato-Rellich Theorem (162 p. [ReS1]), $H = H^0 + q$ is self-adjoint on $\mathfrak{D}(H^0)$ and essentially self-adjoint on any core of H^0. We prove (2.13).

Lemma 2.1. *Let $q \in L^2(0,1)$. Then the estimate (2.13) holds true.*

Proof. For $f \in \mathfrak{D}(H^0), f = (f_\omega)_{\omega\in\mathcal{Z}}, \omega = (n,j,k) \in \mathcal{Z}$, we have the following estimates

$$\|qf_\omega\|_0 \leqslant \max_{x\in\Gamma_\omega} |f_\omega(x)| \|q\|_0, \quad \max_{x\in\Gamma_\omega} |f_\omega(x)| \leqslant \int_{\Gamma_\omega} (|f_\omega| + |f_\omega'|) dx \leqslant \|f_\omega'\|_0 + \|f_\omega\|_0,$$

which yield

$$\|qf\|^2 = \sum \|qf_\omega\|_0^2 \leqslant \|q\|_0^2 \sum (\|f_\omega'\|_0 + \|f_\omega\|_0)^2.$$

Thus for $C_q = \|q\|_0^2$ we obtain

$$\|qf\|^2 \leqslant 2C_q \sum (\|f_\omega'\|_0^2 + \|f_\omega\|_0^2) = 2C_q(\|f'\|^2 + \|f\|^2) \leqslant \varepsilon\|H^0 f\|^2 + \Big(\frac{C_q^2}{\varepsilon} + 2C_q\Big)\|f\|^2,$$

for any $\varepsilon > 0$, since

$$2C_q\|f'\|^2 = 2C_q(H^0 f, f) \leqslant 2C_q\|H^0 f\|\|f\| \leqslant \varepsilon\|H^0 f\|^2 + \frac{C_q^2}{\varepsilon}\|f\|^2,$$

which yields (2.1). ■

Note that the operator H_k in $L^2(\Gamma^1)$ with $\mathfrak{D}(H_k) = \{f, \oplus_{\alpha\in\mathcal{Z}_1} f''_\alpha \in L^2(\Gamma^1) : f \ \ satisfies\ the\ Kirchhoff\ conditions\ (1.4),\ (1.5)\}$ is self-adjoint. The proof is similar to the case H.

Proof of Theorem 1.2. (i) Let $H_kF = \lambda F$ for some eigenfunction F and some $\lambda \in \sigma_D$. For each $F_\omega = F|_{\Gamma_\omega}, \omega = (n, j) \in \mathcal{Z}_1 = \mathbb{Z}\times\mathbb{J}$, we have $F_\omega(t) = F_\omega(0)+\int_0^t F'_\omega(s)ds, t \in [0, 1]$. Then $|F_\omega(0)| \leqslant |F_\omega(x)| + \int_0^1 |F'_\omega(t)|dt$ and integrating over $x \in [0, 1]$, using the Hölder inequality, we obtain

$$|F_\omega(0)| \leqslant \int_0^1 (|F_\omega(t)| + |F'_\omega(t)|)dt \leqslant \|F_\omega\|_0 + \|F'_\omega\|_0 = o(1), \quad as \quad n \to \pm\infty, \tag{2.14}$$

since $F \in \mathcal{H}_k(\lambda)$. Furthermore, each restriction F_ω has the form $F_\omega(t) = a_\omega\vartheta_t+b_\omega\varphi_t, t \in [0, 1]$ for some constants a_ω, b_ω, which implies $F_\omega(1) = F_\omega(0)\vartheta_1$, where $\varphi_t = \varphi(t, \lambda), \vartheta_t = \vartheta(t, \lambda), ..$ The Kirchhoff conditions give $F_{n+1,0}(0) = F_{n,1}(1) = \vartheta_1 F_{n,1}(0) = \vartheta_1 F_{n,0}(1) = \vartheta_1^2 F_{n,0}(0)$, which yields

$$F_{0,0} = \vartheta_1^{2n} F_{-n,0}(0), \quad all\ n \in \mathbb{Z}. \tag{2.15}$$

Let $|\vartheta_1| \leqslant 1$, the proof for the case $|\vartheta_1| > 1$ is similar. Then (2.15), (2.14) imply $F_{0,0} = \vartheta_1^{2n} F_{-n,0}(0) = o(1)$ as $n \to +\infty$. Thus (2.15) gives $F_{n,0}(1) = 0$ for all $n \in \mathbb{Z}$. Finally, F vanishes on all vertexes of Γ^1, since the set of all ends of vertexes $\Gamma_{n,0}, n \in \mathbb{Z}$ coincides with the vertex set of Γ^1.

(ii) Using (1.11)-(1.12), we deduce that $\psi^{(0)}$ satisfies the Kirchhoff conditions (1.4), (1.5). Thus ψ^0 is a eigenfunction of H_k.

The operator H_k is periodic, then each $\psi^{(n)}, n \in \mathbb{Z}$ is an eigenfunction. We will show that the sequence $\psi^{(n)}, n \in \mathbb{Z}$ forms a basis for $\mathcal{H}_k(\lambda)$. The functions $\psi^{(n)}$ are linearly independent, since $\operatorname{supp}\psi^{(n)} \cap \operatorname{supp}\psi^{(m)} = \emptyset$ for all $n \neq m$.

Consider the first case $\eta = 1 - s^k {\varphi'_1}^2 \neq 0$. For any $f \in \mathcal{H}_k(\lambda)$ we will show the identity (1.13), i.e., $f = \widehat{f}$, where $\widehat{f} = \sum_{n\in\mathbb{Z}} \widehat{f}_n\psi^{(n)}$, $\widehat{f}_n = \frac{f'_{n,0}(0)}{\eta}$. The definition of $\widehat{f}$ and $\lambda \in \sigma_D$ give $\widehat{f}|_{\Gamma_{n,0}} = f|_{\Gamma_{n,0}} = \widehat{f}_n\varphi$ for all $n \in \mathbb{Z}$. This yields $\sum |\widehat{f}_n|^2 < \infty$ and $\widehat{f} \in L^2(\Gamma^1)$, since $f \in L^2(\Gamma^1)$

Note that $\widehat{f}$ satisfies the Kirchhoff conditions (1.4), (1.5) and $-\widehat{f}''_\alpha + q\widehat{f}_\alpha = \lambda\widehat{f}_\alpha, \alpha \in \mathcal{Z}_1$. Consider the function $u = f - \widehat{f} = (u_\alpha)_{\alpha\in\mathcal{Z}_1}$, where $u_{n,j} = C_{n,j}\varphi(t, \lambda), j = 1, 2$ and $u_{n,0} = 0$ for some constant $C_{n,j}, n \in \mathbb{Z}$. The Kirchhoff boundary conditions (1.4)-(1.5) yields $u = 0$.

Consider the second case $s^k {\varphi'_1}^2 = 1$. For any $f \in \mathcal{H}_k(\lambda)$ we will show the identity (1.13), i.e., $f = \widehat{f}$, where $\widehat{f} = \sum_{n\in\mathbb{Z}} \widehat{f}_n\psi^{(n)}$, $\widehat{f}_n = f'_{n,1}(0)$. From the definition of $\widehat{f}$ and $\lambda \in \sigma_D$ we deduce that $\widehat{f}|_{\Gamma_{n,1}} = f|_{\Gamma_{n,1}} = \widehat{f}_n\varphi$ for all $n \in \mathbb{Z}$. This yields $\sum |\widehat{f}_n|^2 < \infty$ and $\widehat{f} \in L^2(\Gamma^1)$, since $f \in L^2(\Gamma^1)$.

Consider the function $u = f - \widehat{f}$. The function $u = 0$ at all vertices of Γ^1 and then $u_\alpha = C_\alpha\varphi_t, \alpha = (n, j), n \in \mathbb{Z}, j = 0, 2$. Assume that $C_{0,0} = C$. Then the Kirchhoff boundary conditions (1.5) yields $C_{n,0} = -C_{n,2} = -C$ and $C_{n,0} = C_{n+1,0} = C$, which give $C = 0$ since $u \in L^2(\Gamma^{(1)})$. ■

3 The Lyapunov function Δ_0

Recall that $\vartheta(x,\lambda),\varphi(x,\lambda),x\in\mathbb{R}$ are the fundamental solutions of Eq. $-y''+qy=\lambda y,\ \lambda\in\mathbb{C}$, on the real line such that $\vartheta(0,\lambda)=\varphi'(0,\lambda)=1,\ \vartheta'(0,\lambda)=\varphi(0,\lambda)=1$. For each $x\in\mathbb{R}$ the functions $\vartheta,\vartheta',\varphi,\varphi'$ are entire in $\lambda\in\mathbb{C}$ and satisfy:

$$\vartheta(x,\lambda)=\cos\sqrt{\lambda}x+\frac{1}{2\sqrt{\lambda}}\int_0^x\Big(\sin\sqrt{\lambda}x+\sin\sqrt{\lambda}(x-2t)\Big)q(t)dt+O\Big(\frac{e^{|\operatorname{Im}\sqrt{\lambda}|x}}{|\lambda|}\Big),\tag{3.1}$$

$$\vartheta'(x,\lambda)=-\sqrt{\lambda}\sin\sqrt{\lambda}x+\frac{1}{2}\int_0^x\Big(\cos\sqrt{\lambda}x+\cos\sqrt{\lambda}(x-2t)\Big)q(t)dt+O\Big(\frac{e^{|\operatorname{Im}\sqrt{\lambda}|x}}{|\lambda|^{1/2}}\Big),\tag{3.2}$$

$$\varphi(x,\lambda)=\frac{\sin\sqrt{\lambda}x}{\sqrt{\lambda}}+\frac{1}{2\lambda}\int_0^x\Big(-\cos\sqrt{\lambda}x+\cos\sqrt{\lambda}(x-2t)\Big)q(t)dt+O\Big(\frac{e^{|\operatorname{Im}\sqrt{\lambda}|x}}{|\lambda|^{3/2}}\Big),\tag{3.3}$$

$$\varphi'(x,\lambda)=\cos\sqrt{\lambda}x+\frac{1}{2\sqrt{\lambda}}\int_0^x\Big(\sin\sqrt{\lambda}x-\sin\sqrt{\lambda}(x-2t)\Big)q(t)dt+O\Big(\frac{e^{|\operatorname{Im}\sqrt{\lambda}|x}}{|\lambda|}\Big)\tag{3.4}$$

as $|\lambda|\to\infty$, uniformly on bounded sets of $(x;q)\in[0,1]\times L^2(0,1)$ (see [PT]), and

$$\Delta(\lambda)=\cos\sqrt{\lambda}+\frac{q_0\sin\sqrt{\lambda}}{2\sqrt{\lambda}}+\frac{O(e^{|\operatorname{Im}\sqrt{\lambda}|})}{|\lambda|},\quad q_0=\int_0^1 q(t)dt.\tag{3.5}$$

Substituting (3.1)-(3.5) into Δ_0 we obtain (1.19), i.e.,

$$\Delta_0(\lambda)=\Delta_0^0(\lambda)+\frac{9q_0}{8}\frac{\sin 2\sqrt{\lambda}}{\sqrt{\lambda}}+O\Big(\frac{e^{2|\operatorname{Im}\sqrt{\lambda}|}}{|\lambda|}\Big),\quad \Delta_0^0(\lambda)=\frac{9\cos 2\sqrt{\lambda}-1}{8}.$$

Let $\Delta_-=\frac{1}{2}(\varphi'(1,\cdot)-\vartheta(1,\cdot))$. Substituting the identity $\varphi(1,\cdot)\vartheta'(1,\cdot)=\Delta^2-\Delta_-^2-1$ into $\Delta_0=2\Delta^2(\lambda)+\frac{\vartheta'(1,\lambda)\varphi(1,\lambda)}{4}-1$, we obtain

$$\Delta_0=\frac{9\Delta^2-\Delta_-^2-5}{4},\qquad\text{where}\quad \Delta_-=\frac{\varphi'(1,\cdot)-\vartheta(1,\cdot)}{2}.\tag{3.6}$$

Asymptotics (3.1)-(3.4) yield as $|\lambda|\to\infty$

$$\Delta_-(\lambda)=-\frac{F(\lambda)}{2\sqrt{\lambda}}+\frac{O(e^{|\operatorname{Im}\sqrt{\lambda}|})}{|\lambda|},\qquad F(\lambda)=\int_0^1\sin\sqrt{\lambda}(1-2t)q(t)dt.\tag{3.7}$$

Proof of Theorem 1.3. (i) Using (1.10) we obtain the characteristic equation

$$\det(\mathcal{M}_k-\tau I_2)=\tau^2-2a_k\tau+s^{-k}=0,\qquad a_k=\frac{\operatorname{Tr}\mathcal{M}_k}{2}=\frac{2\xi_k}{1+s^k},\quad \xi_k=\Delta_0+s_k^2.$$

The eigenvalues $\tau_{k,\pm}$ of $\mathcal{M}_k$ are given by $\tau_{k,\pm} = a_k \pm \sqrt{a_k^2 - s^{-k}}$. Using $a_k\frac{(1+s^k)}{2} = \xi_k$ and $\frac{(1-s^k)^2}{4}(a_k^2 - s^{-k}) = \rho_k$ we have

$$\Delta_{k,+} = \frac{1}{2}(\tau_{k,+} + \tau_{k,+}^{-1}) = \frac{1}{2}(\tau_{k,+} + s^k \tau_{k,-}) = a_k \frac{(1+s^k)}{2} + \frac{(1-s^k)}{2}\sqrt{a_k^2 - s^{-k}} = \xi_k + \sqrt{\rho_k}$$

and a similar argument yields $\Delta_{k,-} = \xi_k - \sqrt{\rho_k}$.

(ii) The standard arguments (see [Ca1]) yields $\sigma(H_k) = \sigma_{ac}(H_k) \cup \sigma_{pp}(H_k)$, i.e., $\sigma_{sc}(H_k) = \emptyset$ and $\sigma_{ac}(H_k) = \{\lambda \in \mathbb{R} : \Delta_k(\lambda) \in [-1,1]\}$.

(iii) Let $\Delta_k'(\lambda_0) = 0$ and $\Delta_k(\lambda_0) \in (-1,1)$ for some $\lambda_0 \in \sigma_k, k = 0,..,N$. Then we have the Tailor series $\Delta_k(\lambda) = \Delta_k(\lambda_0) + \zeta^p \frac{\Delta_k^{(p)}(\lambda_0)}{p!} + O(\zeta^{p+1})$ as $\zeta = \lambda - \lambda_0 \to 0$, where $\Delta_k^{(p)}(\lambda_0) \neq 0$ for some $p \geqslant 1$. By the Implicit Function Theorem, there exists some curve $Y \subset \{\lambda : |\lambda - \lambda_0| < \varepsilon\} \cap \mathbb{C}_+, Y \neq \emptyset$, for some $\varepsilon > 0$ such that $\Delta_k(\lambda) \in (-1,1)$ for any $\lambda \in Y$. Thus we have a contradiction with (1.17). ■

In order to prove Theorem 1.4 we need

Lemma 3.1. *Let* $q \in L^2(0,1)$. *Then the following statements hold:*
(i) $\Delta_0(\lambda) \geqslant 1$ *for all* $\lambda = \mu_n$ *and* $\lambda = \nu_{n-1}, n \geqslant 1$. *If in addition* $\Delta^2(\lambda) = 1$, *then* $\Delta_0(\lambda) = 1$.
(ii) If $\Delta(\eta) = 0$ *for some* $\eta \in \mathbb{R}$, *then* $\Delta_0(\eta) = -\frac{5+\vartheta^2(1,\eta)}{4} \leqslant -\frac{5}{4}$.
(iii) Let $q \in L^2_{even}(0,1) = \{q \in L^2_{even}(0,1) : q(x) = q(1-x), x \in [0,1]\}$. *Then*

$$\Delta_0 = \frac{9\Delta^2 - 5}{4}, \qquad \Delta = \varphi'(1,\cdot) = \vartheta(1,\cdot), \tag{3.8}$$

$$\gamma_{0,2n} = \gamma_n, \quad \textit{where each } \gamma_n = (\mu_n, \nu_n), \quad \textit{or} \quad \gamma_n = (\nu_n, \mu_n), \qquad n \geqslant 1. \tag{3.9}$$

The zeros $\eta_n = \lambda_{0,2n-1}$ *for all* $n \geqslant 1$.
(iv) There exists an integer $n_0 > 1$ *such that* $\Delta_0 - 1$ *has exactly* $2n_0 + 1$ *roots, counted with multiplicities, in the domain* $\{\lambda : |\sqrt{\lambda}| < \pi(n_0 + \frac{1}{2})\}$ *and for each* $n > n_0$, *exactly two roots, counted with multiplicities, in the domain* $\{\lambda : |\sqrt{\lambda} - \pi n| < \frac{\pi}{4}\}$. *There are no other roots.*
(v) There exists an integer $n_0 > 1$ *such that* $\Delta_0 + \frac{5}{4}$ *has exactly* $2n_0$ *roots, counted with multiplicities, in the domain* $\{z : |\sqrt{z}| < \pi n_0\}$ *and for each* $n > n_0$, *exactly one simple root in the domain* $\{\lambda : |\sqrt{\lambda} - \pi(n - \frac{1}{2})| < \frac{\pi}{4}\}$. *There are no other roots.*
(vi) There exists an integer $n_0 > 1$ *such that* Δ_0 *has exactly* n_0 *roots, counted with multiplicities, in the domain* $\{\lambda : |\sqrt{\lambda}| < \pi n_0\}$ *and for each* $n > n_0$, *exactly one simple root in the domain* $\{\lambda : |\sqrt{\lambda} - \pi(n - \frac{1}{2})| < \frac{\pi}{4}\}$. *There are no other roots.*
(vii) Let $c \in (-\frac{5}{4}, 1)$ *and let* $u_n^{\pm} = \pi n \pm u_0^+, u_0^+ = \frac{\arccos \frac{1+8c}{9}}{2} \in [0, \frac{\pi}{2}]$, $n \geqslant 1$ *and* $u_0^+ < u_1^- < u_1^+ < u_2^- < ...$ *Then there exists an integer* $n_0 > 1$ *such that* $\Delta_0 - c$ *has exactly* $2n_0 + 1$ *roots, counted with multiplicities, in the domain* $\{\lambda : |\sqrt{\lambda}| < R\}, R = (u_{n_0}^+ + u_{n_0+1}^-)/2$ *and for each* $n > n_0$, *exactly one simple root in the domain* $\{\lambda : |\sqrt{\lambda} - u_n^{\pm}| < r$ *for some small* $r \in (0,1)$. *There are no other roots.*

Proof. (i) Let $\lambda \in \{\mu_n, \nu_n\}$ for some $n \geqslant 1$. Then we have $\Delta_0(\lambda) = 2\Delta^2(\lambda) + \frac{\vartheta'(1,\lambda)\varphi(1,\lambda)}{4} - 1 = 2\Delta^2(\lambda) - 1 \geqslant 1$. Moreover, if $\lambda \in \{\mu_n, \nu_n\}$ and $\Delta^2(\lambda) = 1$, then we have $\Delta_0(\lambda) = 1$.

(ii) We have $0 = 2\Delta(\eta) = \varphi'(1,\eta) + \vartheta(1,\eta)$, which yields $\varphi'(1,\eta) = -\vartheta(1,\eta)$. Substituting the last identity into $\Delta_0(\lambda) = 2\Delta^2(\lambda) + \frac{\vartheta'\varphi}{4} - 1$ we get the needed estimate.

(iii) It is well known that if $q \in L^2_{even}(0,1)$, then $\Delta = \varphi'(1,\cdot) = \vartheta(1,\cdot)$, see p.8, [MW]. It is well known that if $\Delta = \varphi'(1,\cdot) = \vartheta(1,\cdot)$, then $q \in L^2_{even}(0,1)$, see [PT].

(iv) Define the contours $C_n(r) = \{\lambda \in \mathbb{C} : |\sqrt{\lambda} - \pi n| = \pi r\}$. Fix another integer $n_1 > n_0$. Consider the contours $C_0(n_0 + (1/2)),\ C_0(n_1 + (1/2))),\ C_n(1/4), n > n_0$. Using the estimate $e^{|\operatorname{Im}\sqrt{\lambda}|} < 4|\sin\sqrt{\lambda}|, |\lambda - \pi n| \geqslant \frac{\pi}{4}$ and (1.19) we obtain on all contours (for large n_0)

$$|(\Delta_0(\lambda) - 1) - (\Delta_0^0(\lambda) - 1)| = o(e^{2|\operatorname{Im}\sqrt{\lambda}|}) = |\sin\sqrt{\lambda}|^2 o(1) = o(1)|\Delta_0^0(\lambda) - 1|$$

Hence, by Rouché's Theorem, $\Delta_0(\lambda) - 1$ has as many roots, counted with multiplicities, as $\Delta_0^0(\lambda) - 1$ in each of the bounded domains and the remaining unbounded domain. Since $\Delta_0^0(\lambda) - 1$ has exactly the simple root $= 0$ and one root of the multiplicity 2 at each $\pi^2 n^2 \geqslant 1$, and since $n_1 > n_0$ can be chosen arbitrarily large, the point (iii) follows.

The proof of (v) and (vi) is similar. ■

Lemma 3.2. *Let $c \in [-\frac{5}{4}, 1]$. Then the equation $\Delta_0(\lambda) = c, \lambda \in \mathbb{C}$ has only real zeros, which satisfy*
(i) If $c \in (-\frac{5}{4}, 1)$ and $q_0 = \int_0^1 q(t)dt$, then these zeros $z_n^\pm$ are given by

$$z_0^+ < z_1^- < z_1^+ < z_2^- < .. \quad \text{and} \quad \sqrt{z_n^\pm} = u_n^\pm + \frac{q_0}{2u_n^\pm} + \frac{o(1)}{n^2} \quad \text{as} \quad n \to \infty, \tag{3.10}$$

where $u_n^\pm = \pi n \pm u_0^+,\ n \geqslant 1$ and $u_0^+ = \frac{\arccos\frac{1+8c}{9}}{2} \in [0, \frac{\pi}{2}], u_0^+ < u_1^- < u_1^+ < u_2^- < ...$

Moreover, the zeros $z_n^\pm$ have another forms given by: $x_n^- = z_{n+1}^+, x_n^+ = z_n^-$, and

$$x_1^- < x_1^+ < x_2^- < x_2^+ < x_3^- < .. \qquad \sqrt{x_n^\pm} = v_n^\pm + \frac{q_0}{2v_n^\pm} + \frac{o(1)}{n^2} \quad \text{as} \quad n \to \infty, \tag{3.11}$$

where $v_n^\pm = \pi(n - \frac{1}{2}) \pm (\frac{\pi}{2} - u_0^+)$.
(ii) If $c = 1$ or $c = -\frac{5}{4}$. Then these zeros $z_n^\pm$ and $x_n^\pm$ are given by

$$z_0^+ < z_1^- \leqslant z_1^+ < z_2^- \leqslant z_2^+ < .. \quad \text{and} \quad \sqrt{z_n^\pm} = \pi n + \frac{q_0}{2\pi n} + \frac{o(1)}{n^2}, \quad if \quad c = 1,$$

$$x_1^- \leqslant x_1^+ < x_2^- \leqslant x_2^+ < .. \quad \text{and} \quad \sqrt{x_n^\pm} = \pi(n - \frac{1}{2}) + \frac{q_0}{2\pi n} + \frac{o(1)}{n^2}, \quad if \quad c = -\frac{5}{4},$$

as $n \to \infty$.

Proof. (i) Let $z = \sqrt{\lambda}$. Recall the folowing asymptotics from [KK1]:

$$\Delta(\lambda) = \cos\chi(\lambda), \quad \chi(\lambda) = \sqrt{\lambda} - \frac{q_0}{2\sqrt{\lambda}} + \frac{o(1)}{\lambda} \qquad as \quad |\lambda| \to \infty. \tag{3.12}$$

Using (3.6) we rewrite the Eq. $\Delta_0(\lambda) = c, \lambda \in \mathbb{R}$ in the form $c = \frac{9\Delta^2 - \Delta_-^2 - 5}{4}$, which yields

$$\cos 2\chi(\lambda) = A_c + \frac{2\Delta_-^2(\lambda)}{9}, \quad A_c = \frac{1 + 8c}{9} \in (-1, 1). \tag{3.13}$$

In the case $q = 0$ we have $\cos 2z = A_c$ and the corresponding solutions $\sqrt{z_n^\pm} = u_n^\pm$. Then by Lemma 3.1, $|\sqrt{z_n^\pm} - u_n^\pm| < r$ for all $n > n_0$, for some $n_0 \geqslant 1$ and some small $r \in (0, 1)$. Using (3.13),(3.7) we obtain $\cos 2\chi(\lambda) = A_c + o(\lambda^{-1})$ as $\lambda \to \infty$, and then $\zeta_n^\pm = \sqrt{z_n^\pm} - u_n^\pm \to 0$. Thus $\varepsilon_n^\pm = \zeta_n^\pm - \frac{q_0}{2u_n^\pm} + \frac{o(1)}{n^2} \to 0$ and the Taylor series gives

$$\cos 2\chi(\lambda) = \cos 2(u_n^\pm + \varepsilon_n^\pm) = \cos 2u_n^\pm - \sin(2u_n^\pm)\varepsilon_n^\pm(1 + O(\varepsilon_n^\pm)) = A_c + o(n^{-2}).$$

Thus we obtain $\varepsilon_n^\pm = \zeta_n^\pm - \frac{2_0}{2u_n^\pm} + \frac{o(1)}{n^2} = \frac{o(1)}{z^2}$ which implies, $\zeta_n^\pm = \frac{q_0}{2u_n^\pm} + \frac{o(1)}{n^2}$.

Proof of (3.11) and (ii) is similar. ■

Proof of Theorem 1.4. (i) The function $\Delta_0 = 2\Delta^2 + \frac{\vartheta'(1,\cdot)\varphi(1,\cdot)}{4} - 1$ is entire and real on real line. Moreover, it has asymptotics (1.19). By Theorem 1.3, the function Δ_0 has only simple real zeros $\eta_{0,n}, n \geqslant 1$, which satisfy $\eta_{0,1} < \eta_{0,2} < \eta_{0,3} < ....$ Then by the Laguerre Theorem (see Sect. 8.52 [Ti]), the function Δ_0' has only real simple zeros $\lambda_{0,n}, n \geqslant 1$, which are separated by the zeros of Δ_0: $\eta_{0,1} < \lambda_{0,1} < \eta_{0,2} < \lambda_{0,2} < \eta_{0,3} < ...$, since $\Delta_0(\lambda) \to \infty$ as $\lambda \to -\infty$. Moreover, we obtain $(-1)^n\Delta_0(\lambda_{0,n}) \geqslant 1, n \geqslant 1$.

Define an interval $\mathcal{G} = (-\infty, \pi^2 n_0^2)$. Lemma 3.1 gives for large integer $n_0 > 1$:

A) The function $\Delta_0 - 1$ has $2n_0 + 1$ zeros on $\mathcal{G}$.

B) $\Delta_0(\eta_{0,n}) \leqslant -\frac{5}{4}, n = 1, .., n_0$, where $\eta_n \in \mathcal{G}$ is a zero of Δ, $\eta_1 < \eta_2 < ....$

C) $\Delta_0(\mu_n) \geqslant 1, n = 1, .., n_0$, where $\mu_n \in \mathcal{G}$ are zeros of $\varphi(1, \lambda)$, the Dirichlet eigenvalues.

These facts and $\mu_1 < \eta_1 < \mu_2 < \eta_2 < ...$ yield $\mu_n \in \widetilde{\gamma}_{0,2n}$ and $\eta_n \in \widetilde{\gamma}_{0,2n-1}$ for all $n \geqslant 1$, where $\widetilde{\gamma}_{0,n} = [\lambda_{0,n}^-, \lambda_{0,n}^+]$. Moreover, we obtain (1.20).

(ii) Using (i) we deduce that the periodic and anti-periodic eigenvalues $\lambda_{0,n}^\pm, n \geqslant 0$ satisfy (1.18). Consider $\lambda_{0,2n+1}^\pm$, which satisfies $\Delta_0(\lambda) = -1$. Using Lemma 3.2 (i) for $c = -1$ we have the first asymptotics in (1.21). In order to show (1.21) we consider $\lambda_{0,2n}^\pm$ with even $n \geqslant 1$. The proof for odd n is similar. Using (3.1)-(3.4) and (3.7) we obtain

$$\dot{\Delta}_0(\lambda) = \frac{O(1)}{n^2}, \quad \ddot{\Delta}_0(\lambda) = -\frac{1 + \frac{O(1)}{n}}{(\pi n)^2}, \quad \Delta_-(\lambda) = \frac{q_{sn} + \frac{O(1)}{n}}{2\pi n}, \quad \dot{\Delta}_-(\lambda) = -\frac{\widetilde{q}_{cn} + \frac{O(1)}{n}}{(2\pi n)^2}, \tag{3.14}$$

$$\dot{\Delta}(\lambda) = \frac{O(1)}{n^2}, \ \ddot{\Delta}(\lambda) = -\frac{1 + \frac{O(1)}{n}}{(\pi n)^2}, \quad \dddot{\Delta}(\lambda) = \frac{O(1)}{n^4}, \quad where \quad \dot{\Delta} = \frac{\partial \Delta}{\partial \lambda} \tag{3.15}$$

as $\sqrt{\lambda} = \pi n + O(1/n)$, where $\widetilde{q}_{cn} = \int_0^1 (1 - 2t)q(t) \cos 2\pi n t dt$. Using $\dot{\Delta}_0(\lambda_{0,2n}) = 0$, we have

$$\dot{\Delta}_0(\lambda_n) = \ddot{\Delta}_0(\lambda_{0,2n}) s_n(1 + O(s_n)), \quad s_n = \lambda_n - \lambda_{0,2n} \quad as \quad n \to \infty.$$

The identity $\dot{\Delta}(\lambda_n) = 0$ and $\Delta_0 = \frac{9\Delta^2 - \Delta_-^2 - 5}{4}$ and (3.14) yield

$$\dot{\Delta}_0(\lambda_n) = -\frac{\Delta_-(\lambda_n)\dot{\Delta}_-(\lambda_n)}{2} = \frac{(q_{sn} + O(\frac{1}{n}))(\widetilde{q}_{cn} + O(\frac{1}{n}))}{(2\pi n)^3}. \tag{3.16}$$

Thus asymptotics (3.14)-(3.16) give

$$s_n = \lambda_n - \lambda_{0,2n} = -\frac{(q_{sn} + O(\frac{1}{n}))(\widetilde{q}_{cn} + O(\frac{1}{n}))}{8\pi n}, \tag{3.17}$$

which implies that roughly speaking the point $\lambda_{0,2n}$ is in the center of the gap γ_n, since $\lambda_n = (\pi n)^2 + q_0 + O(1/n)$ see [Ko].

We will determine asymptotics of $\gamma_{0,2n} = (\lambda_{0,2n}^-, \lambda_{0,2n}^+)$. Due to (3.15) we get

$$\Delta(\lambda_{0,2n}^\pm) = \Delta(\lambda_n) + A_{0,n}^\pm, \quad A_{0,n}^\pm = \ddot{\Delta}(\lambda_n)\frac{{\zeta_{0,2n}^\pm}^2}{2}\Big(1 + \frac{O(\zeta_{0,2n}^\pm)}{n^2}\Big), \quad \zeta_{0,2n}^\pm = \lambda_{0,2n}^\pm - \lambda_n, \tag{3.18}$$

$$1 = \Delta(\lambda_n^\pm) = \Delta(\lambda_n) + A_n^\pm, \quad A_n^\pm = \ddot{\Delta}(\lambda_n)\frac{\zeta_n^{\pm 2}}{2}\Big(1 + \frac{O(\zeta_n^\pm)}{n^2}\Big), \quad \zeta_n^\pm = \lambda_n^\pm - \lambda_n, \tag{3.19}$$

which yields $\Delta(\lambda_{0,n}^\pm) - 1 = -A_n^\pm + A_{0,n}^\pm$. Substituting (3.14) into the identity $\Delta(\lambda_{0,n}^\pm) - 1 = \sqrt{1 + \frac{\Delta_-^2(\lambda_{0,n}^\pm)}{9}} - 1$ we obtain

$$\Delta(\lambda_{0,n}^\pm) - 1 = \sqrt{1 + \frac{\Delta_-^2(\lambda_{0,n}^\pm)}{9}} - 1 = \frac{(q_{sn} + O(n^{-1}))^2}{18(2\pi n)^2}. \tag{3.20}$$

Substituting (3.20) and (3.14) into the identity $\Delta(\lambda_{0,n}^\pm) - 1 = -A_n^\pm + A_{0,n}^\pm$ we get

$$\frac{\hat{q}_{sn}^2}{9} + \frac{O(|\hat{q}_n|)}{n} = \zeta_n^{\pm 2} - {\zeta_{0,n}^\pm}^2 + \frac{O(1)}{n^2}.$$

Using $\lambda_n^\pm = (\pi n)^2 + q_0 \pm |\hat{q}_n| + O(n^{-1})$ from [MO] and $\lambda_n = (\pi n)^2 + q_0 + O(n^{-1})$ from [Ko] we obtain $\zeta_n^\pm = \lambda_n^\pm - \lambda_n = \pm|\hat{q}_n| + O(n^{-1})$ which yields

$$\zeta_{0,n}^\pm = \pm\sqrt{\zeta_n^{\pm 2} - \frac{\hat{q}_{sn}^2}{9} + \frac{O(|\hat{q}_n|)}{n}} = \pm\sqrt{|\hat{q}_n|^2 - \frac{\hat{q}_{sn}^2}{9}} + O(n^{-1}). \tag{3.21}$$

(iii) Let μ_n, ν_n lay on the different edge of the gap γ_n. Then Lemma 3.1 (i) gives $\gamma_n = \gamma_{0,2n}$.

Conversely, let $\gamma_n = \gamma_{0,2n} = (a,b)$ and let $\lambda \in \{a,b\}$. Then the identity $\Delta_0 = 2\Delta^2 + \frac{\vartheta'(1,\cdot)\varphi(1,\cdot)}{4} - 1$ yields $1 = \Delta_0(\lambda) = 1 + \frac{\vartheta'(\lambda)\varphi(\lambda)}{4}$, thus $\vartheta'(\lambda)\varphi(\lambda) = 0$, which gives (iii)

(iv) Let $\gamma_n = (a,b)$ and let $a < \mu_n < \nu_n < b$. The proof of other cases is similar. Let $\Delta_0 = f_0 + f$, where $f_0 = 2\Delta^2 - 1, f = \frac{\vartheta'(1,\lambda)\varphi(1,\lambda)}{4}$. We obtain

$$f_0|_{\gamma_n} > 1, \qquad f|_\sigma > 0, \qquad f|_{\gamma_n \setminus \sigma} < 0 \quad \textit{where} \quad \sigma = (\mu_n, \nu_n),$$

which yields $\Delta_0|_\sigma > 1$ and $\Delta_0(\lambda) < 1$ for any $\lambda \in \{a,b\}$. Then there exist two points $\lambda_n^\pm \in (a,b)$ such that $\Delta_0(\lambda_n^\pm) = 1$. Thus we have $\gamma_{0,2n} \subset \gamma_n$ for all $n \geqslant 1$.

We will show $\gamma_{0,2n} = \emptyset$ iff $\gamma_n = \emptyset$. If $\gamma_n = \emptyset$, then $\gamma_{0,2n} \subset \gamma_n$ for all $n \geqslant 1$ yields $\gamma_{0,2n} = \emptyset$.

If $\gamma_{0,2n} = \emptyset$, then for $\lambda = \{\mu_n, \nu_n\}, n \geqslant 1$ we obtain $\Delta_0(\lambda) = 2\Delta^2(\lambda) + \frac{\vartheta'(1,\lambda)\varphi(1,\lambda)}{4} - 1 = 2\Delta^2(\lambda) - 1 = 0$, which yields $\Delta^2(\lambda) = 1$. Thus we deduce that $\gamma_{0,2n} = \emptyset$ iff $\gamma_n = \emptyset$.

(v) Let $q \in L^2_{even}(0,1)$. Then we get $\varphi'(1,\cdot) = \vartheta(1,\cdot) = \Delta$ (see p. 8, [23]), which together with (3.6) yields $\Delta_0 = \frac{9}{4}\Delta^2 - \frac{5}{4}$. Then the zeros of $\Delta(\lambda)$ and $\Delta_0'(\lambda)$(at $\Delta_0(\lambda) < 0$) coincide, since all zeros of Δ_0' are simple.

Conversely, let $\Delta_0(\lambda_{0,n}) = -\frac{5}{4}$ for all odd $n \geqslant 1$. Then Lemma 3.1 ii) gives $\vartheta(1,\lambda_{0,n}) = \Delta(\lambda_{0,n}) = \varphi'(1,\lambda_{0,n}) = 0$ for all odd $n \geqslant 1$. Then $\eta_n = \eta_{0,2n-1}, n \geqslant 1$, which implies $\varphi'(1,\cdot) = \vartheta(1,\cdot) = \Delta$. Thus the Wronskian identity gives $\Delta^2 - \vartheta'(1,\cdot)\varphi(1,\cdot) = 1$ and all zeros of $\Delta^2(\lambda) = 1$ coincide with the Dirichlet and Neumann eigenvalues. Then the results of [GT] or [KK] imply $q \in L^2_{even}(0,1)$. It is simple fact in the inverse spectral theory and it can be proved using other methods, see [PT]. ■

4 The Lyapunov function $\Delta_k, k = 1,..,m$

Proof of Theorem 1.5. (i) We determine the equation for periodic eigenvalues. Using (1.10) and $4c_k^2 = (1+s^k)(1+s^{-k})$ for $k \in \overline{m} = \{1,2,..,m\}$, we have

$$\det(\mathcal{M}_k \mp I_2) = 1 \mp \operatorname{Tr} M_k + s^{-k} = 1 + s^{-k} \mp 4\frac{\Delta_0 + s_k^2}{(1+s^k)} = \frac{4(c_k^2 \mp (\Delta_0 + s_k^2))}{(1+s^k)},$$

which yields the equation

$$\Delta_0(\lambda) = c_k^2 - s_k^2 = \cos\frac{2\pi k}{N} \in (-1,1) \setminus \{0\} \quad \text{for periodic eigenvalues} \ \ \lambda, \tag{4.1}$$

$$\Delta_0(\lambda) = -c_k^2 - s_k^2 = -1 \qquad \text{for anti}-\text{periodic eigenvalues} \ \ \lambda. \tag{4.2}$$

Then by Lemma 3.2, all zeros of $\det(\mathcal{M}_k \mp I_2) = 0$ are real and simple. Using $\xi_k = \Delta_0 + s_k^2, \quad \rho_k = \frac{s_k^2}{c_k^2}(c_k^2 - \xi_k^2)$, we get (recall $\dot u = \frac{\partial}{\partial\lambda}u$)

$$\Delta_k = \xi_k + \sqrt{\rho_k} = \Delta_0 + s_k^2 + \sqrt{\rho_k}, \qquad \dot\Delta_k = \dot\xi_k + \frac{\dot\rho_k}{2\sqrt{\rho_k}} = \left(1 - \frac{s_k^2}{c_k^2}\frac{\xi_k}{\sqrt{\rho_k}}\right)\dot\Delta_0. \tag{4.3}$$

For periodic eigenvalues $\lambda = \lambda_{k,n}^\pm$, where $n \geqslant 0$ is even, we have $\Delta_k(\lambda) = 1$ and

$$\xi_k = \Delta_0 + s_k^2 = c_k^2 - s_k^2 + s_k^2 = c_k^2, \qquad \sqrt{\rho_k} = 1 - \xi_k = 1 - c_k^2 = s_k^2, \ \ at \ \lambda = \lambda_{k,n}^\pm, \tag{4.4}$$

and then (4.3) yields at $\lambda = \lambda_{k,n}^\pm$:

$$\frac{\dot\Delta_k}{\dot\Delta_0} = 1 - \frac{s_k^2}{c_k^2}\frac{\xi_k}{\sqrt{\rho_k}} = 1 - 1 = 0, \quad \ddot\Delta_k = -\frac{s_k^2\dot\Delta_0}{c_k^2}\left(\frac{\dot\Delta_0}{\sqrt{\rho_k}} - \frac{\xi_k\dot\rho_k}{2\rho_k^{3/2}}\right) = -\frac{\dot\Delta_0^2}{c_k^2}\left(1 + \frac{c_k^2}{s_k^2}\right) < 0,$$

which gives (1.23) for periodic eigenvalues $\lambda_{k,2n}^\pm$.

For anti-periodic eigenvalues $\lambda = \lambda_{k,n}^\pm$, where $n \geqslant 1$ is odd, we nave $\Delta_k(\lambda) = -1$ and

$$\xi_k = \Delta_0 + s_k^2 = -1 + s_k^2 = -c_k^2, \qquad \sqrt{\rho_k} = -1 - \xi_k = -s_k^2, \ \ at \ \lambda = \lambda_{k,n}^\pm, \tag{4.5}$$

and then (4.3) yields

$$\frac{\dot\Delta_k}{\dot\Delta_0} = 1 - \frac{(1-c_k^2)}{c_k^2}\frac{(1-s_k^2)}{s_k^2} = 0, \quad \ddot\Delta_k = -\frac{s_k^2\dot\Delta_0}{c_k^2}\left(\frac{\dot\Delta_0}{\sqrt{\rho_k}} - \frac{\xi_k\dot\rho_k}{2\rho_k^{3/2}}\right) = \frac{\dot\Delta_0^2}{c_k^2}\left(1 + \frac{c_k^2}{s_k^2}\right) > 0,$$

which yields (1.24) for anti-periodic eigenvalues $\lambda_{k,n}^{\pm}$.

Let $\lambda = \lambda_{k,2n}^{\pm}, k \in \overline{m}, n \geqslant 0$, be periodic eigenvalues. Recall that they satisfy $\Delta_0(\lambda) = \cos\frac{2\pi k}{N} \in (-1,1)$. Using Lemma 3.2 for $c = \cos\frac{2\pi k}{N}$ we have (1.25).

(ii) We determine the equation for resonances for $k \in \overline{m}$. We have: $\rho_k(\lambda) = 0 \Leftrightarrow \xi_k(\lambda) = \pm c_k$. Consider the first case $+$: n is **even**, i.e., $\frac{n}{2} \in \mathbb{Z}$. We obtain

$$\xi_k(\lambda) = c_k \qquad \Leftrightarrow \quad \Delta_0(\lambda) > a_k^+ = c_k - s_k^2 \in (-1,1), \quad k \in \overline{m}. \tag{4.6}$$

Then the resonances $r_{k,n}^{\pm}$ are zeros of Eq. $\Delta_0(\lambda) = c_k - s_k^2 \in (-1,1), \lambda \in \mathbb{C}$. From Lemma 3.2 we deduce that all these resonances are real. Moreover, we have $1 > a_1^+ > a_2^+ > ... > a_m^+$, which yields for even all even $n \geqslant 1$:

$$r_{1,0}^+ < r_{2,0}^+ < ... < r_{m,0}^+, \quad r_{m,n}^- ... < r_{2,n}^- < r_{1,n}^- < \lambda_{0,n}^- < \lambda_{0,n}^+ < r_{1,n}^+ < r_{2,n}^+ < ... < r_{m,n}^+,$$

$$\gamma_{m,n} \subset \gamma_{m-1,n} \subset ... \subset \gamma_{1,n} \subset \gamma_{0,n}, \qquad \textit{and} \;\; G_n = \cap_0^m \gamma_{k,n} = \gamma_{0,n}. \tag{4.7}$$

Consider the second case $-$: n is **odd**. We have

$$\xi_k(\lambda) = -c_k \qquad \Leftrightarrow \quad \Delta_0(\lambda) = a_k^- = -c_k - s_k^2 \in (-\frac{5}{4}, -1), \quad \frac{N}{3} \notin \mathbb{Z}, \tag{4.8}$$

$$\xi_k(\lambda) = -c_k \qquad \Leftrightarrow \quad \Delta_0(\lambda) = a_k^- = -c_k - s_k^2 \in [-\frac{5}{4}, -1), \quad \frac{N}{3} \in \mathbb{Z}. \tag{4.9}$$

Then from Theorem 1.3 we deduce that the zeros of ρ_k are zeros of Eq. $\Delta_0(\lambda) = -c_k - s_k^2 \in [-\frac{5}{4}, -1), \lambda \in \mathbb{C}$. Moreover, by Lemma 3.2, they are real and simple.

If $\frac{N}{3} \notin \mathbb{Z}$, then $\frac{N}{3} = p + \varepsilon$ for some integer $p \geqslant 1$ and $\varepsilon \in (0,1)$. Then we obtain

$$\gamma_{0,n} \supset \gamma_{1,n} \supset \gamma_{2,n} \supset ... \supset \gamma_{p,n}, \quad \textit{and} \quad \gamma_{p+1,n} \subset \gamma_{p+2,n} \subset .. \subset \gamma_{m,n}, \tag{4.10}$$

$$\gamma_{p,n} \subset \gamma_{p+1,n} \;\; \textit{or} \quad \gamma_{p,n} \supset \gamma_{p+1,n} \;\; \textit{and} \;\; G_n = \cap_1^N \gamma_{k,n} = \gamma_{p,n} \cap \gamma_{p+1,n}, \tag{4.11}$$

for all odd $n \geqslant 1$. If $p = \frac{N}{3} \in \mathbb{Z}$, then we get for all odd $n \geqslant 1$:

$$\gamma_{0,n} \supset \gamma_{1,n} \supset \gamma_{2,n} \supset ... \supset \gamma_{p,n}, \quad \textit{and} \quad \gamma_{p,n} \subset \gamma_{p+1,n} \subset .. \subset \gamma_{m,n}, \tag{4.12}$$

$$G_n = \cap_{k=0}^m \gamma_{k,n} = \gamma_{p,n}, \quad \Delta_0(r_{p,n}^{\pm}) = -\frac{5}{4}. \tag{4.13}$$

Resonances $r_{k,2n}^{\pm}, k \in \overline{m}, n \geqslant 0$ satisfies $\Delta_0(r_{k,2n}^{\pm}) = c_k - s_k^2 \in (-1,1)$. Using Lemma 3.2 for $c = c_k - s_k^2$ we obtain asymptotics (1.29) for $r_{k,2n}^{\pm}$.

Resonances $r_{k,2n+1}^{\pm}, k \in \overline{m}, k \neq \frac{N}{3}, n \geqslant 0$ satisfies $\Delta_0(r_{k,2n+1}^{\pm}) = -c_k - s_k^2 \in (-\frac{5}{4}, -1)$. Using Lemma 3.2 for $c = c_k - s_k^2$ we obtain asymptotics (1.29) for $r_{k,2n+1}^{\pm}$.

(iii) Finally, we obtain the following identities for $(n,k) \in \mathbb{N} \times \overline{m}$:

$$S_n = \cup_{k=0}^m \sigma_{k,n}, \quad \sigma_{0,n} = (\lambda_{0,n-1}^+, \lambda_{0,n}^-), \quad \sigma_{k,n} = (r_{k,n-1}^+, r_{k,n}^-), \tag{4.14}$$

$$G_n = \cap_{k=0}^m \gamma_{k,n}, \ \gamma_{0,n} = (\lambda_{0,n}^-, \lambda_{0,n}^+), \qquad \gamma_{k,n} = (r_{k,n}^-, r_{k,n}^+), \tag{4.15}$$

which together with (4.7), (4.10)-(4.13) yield gives (1.30)-(1.32).

(iv) Using Theorem 1.4 we deduce that $G_{2n} = \gamma_{0,2n} = \emptyset$ iff $\gamma_n = \emptyset$. Moreover, asymptotics $|\gamma_n| \to 0$ as $n \to \infty$ give $|G_{2n}| \to 0$ as $n \to \infty$.

(v) In the case $p = \frac{N}{3} \in \mathbb{Z}$ and odd $n \geqslant 1$ using the identity (4.13), we have $G_n = \gamma_{p,n} = (r_{p,n}^-, r_{p,n}^+)$, where $r_{p,n}^\pm$ are zeros of Eq. $\Delta_0(\lambda) = -\frac{5}{4}$. Theorem 1.4 (v) gives $r_{p,n}^\pm = \lambda_n$ iff $q \in L^2_{even}(0,1)$.

We determine the asymptotics (1.33). Let $(z_n^\pm)^2 = r_{p,n}^\pm$. Using $\frac{9\Delta^2 - \Delta_-^2 - 5}{4} = \frac{-5}{4}$ we have $9\Delta^2(r_{p,n}^\pm) = \Delta_-^2(r_{p,n}^\pm)$. Thus in the case $q = 0$ we have $\cos z = 0$ and the corresponding zeros are given by $z_n = \pi(n - \frac{1}{2}), n \geqslant 1$.

In the case $q \neq 0$ we have $\Delta(\lambda) = \pm\frac{1}{3}\Delta_-(\lambda) = \frac{o(1)}{\sqrt{\lambda}}$ as $\lambda \to \infty$. Then by Lemma 3.1, $|z_n^\pm - z_n| < \frac{\pi}{4}$ for all $n > n_0$, for some $n_0 \geqslant 1$. Thus we obtain $z_n^\pm = z_n + \zeta_n^\pm, \zeta_n^\pm \to 0$. Moreover, using (3.12), we get $\frac{o(1)}{n} = \Delta(z_n + \zeta_n^\pm) = \cos(z_n + \zeta_n^\pm - \frac{q_0}{2z_n} + \frac{o(1)}{n^2})$, which implies $\zeta_n^\pm = \frac{q_0}{2z_n} + \frac{o(1)}{n}$. Let $\zeta_n^\pm = \frac{q_0 + v}{2z_n}, v_n^\pm \to 0$ and for $R_n^\pm = \frac{v_n^\pm}{2z_n} + \frac{o(1)}{n^2}$ using (3.12) we obtain

$$\Delta(z_n + \zeta_n^\pm) = \cos(z_n + R_n^\pm) = (-1)^{n-1}\cos(-\frac{\pi}{2} + R_n^\pm) = (-1)^{n-1}\sin(\frac{v_n^\pm}{2z_n} + \frac{o(1)}{n^2})$$

Using this and (3.7) we get

$$\Delta_-(r_{p,n}^\pm) = (-1)^n \frac{\widetilde{q}_{cn}}{2z_n} + \frac{O(1)}{n^2}, \qquad n \to \infty. \tag{4.16}$$

Thus the Eq. $\Delta(r_{p,n}^\pm) = \pm\frac{1}{3}\Delta_-(r_{p,n}^\pm)$ yields (1.33).

(vi) In the case $\frac{N}{3} = p + \varepsilon$, where $p \geqslant 1$ is integer and $\varepsilon \in (0,1)$ and odd $n \geqslant 1$ the identity (4.11) we have $G_n = \gamma_{p,n} \cap \gamma_{p+1,n}$. In (1.29) we will show that $|\gamma_{k,n}| \to \infty$ as $n \to \infty, k \neq \frac{N}{3}$, which yields $|G_n| \to \infty$ as $n \to \infty$. ■

We will prove the last Corollary 1.6.

Proof of Corollary 1.6. (i) Recall that by Theorem 1.5, each $G_{2n} = \gamma_{0,2n} \subset \gamma_n, n \geqslant 1$. Thus by Theorem 1.5 (iv), $G_{2n} = \gamma_{0,2n} = \emptyset$ for all $n \geqslant n_0$ and for some $n_0 \geqslant 1$ iff q is a finite gap potential for the operator $-y'' + qy$ on the real line.

Consider the odd gaps $G_{2n-1}, n \geqslant 1$. Recall that by Theorem 1.5, if $\frac{N}{3} \notin \mathbb{N}$, then $|G_{2n-1}| \to \infty$ as $n \to \infty$.

Assume that $p = \frac{N}{3} \in \mathbb{N}$. In this case by Theorem 1.5, $G_{2n+1} = \gamma_{p,2n+1} = \emptyset$ iff $q \in L^2_{even}(0,1)$. Thus the statement i) has been proved.

(ii) The function f_ω satisfies the Eq. $-f_\omega'' + qf_\omega = E_n f_\omega$ on the interval $[0,1]$ and $f_\omega(0) = f_\omega(1) = 0, f_\omega'(1)^2 = e^{2h_n} > 0$ for some $h_n \in \mathbb{R}$. Each constant h_n is so-called norming constant for the Sturm-Liouville problem $-y'' + qy = \lambda y$ on the interval $[0,1]$ with the Dirichlet boundary conditions $y(0) = y(1) = 0$ [PT].

Recall that $\mu_n, n \geqslant 1$, is the Dirichlet spectrum of the problem $-y'' + qy = \lambda y, y(0) = y(1) = 0$ on the unit interval $[0,1]$ and $\mu_n = E_n, n \geqslant 1$.

Recall the well know result from [PT]: the mapping $\mathfrak{F} : q \to \Big(q_0, (\varkappa_n(q))_1^\infty; (h_n(q))_1^\infty \Big)$ is a real-analytic isomorphism between $L^2(0,1)$ and $\mathbb{R} \times \mathcal{K} \times \ell_1^2$. This gives the statement ii).

(iii) If potential q is even, i.e., $q \in L^2_{even}(0,1)$, then each $h_n = 0, n \geqslant 1$ (see [PT]).

Recall the well know result from [PT]: the mapping $\mathfrak{F}_e : q \to \Big(q_0, (\varkappa_n(q))_1^\infty \Big)$ is a real-analytic isomorphism between $L^2_{even}(0,1)$ and $\mathbb{R} \times \mathcal{K}$. This gives the statement iii). ■

Acknowledgments. E. Korotyaev was partly supported by DFG project BR691/23-1. The various parts of this paper were written at the Mittag-Leffler Institute, Stockholm and in the Erwin Schrödinger Institute for Mathematical Physics, Vienna, the first author is grateful to the Institutes for the hospitality. The first author would like to thank Markus Klein for useful discussions.

References

[ARZ] Avron, J. E.; Raveh, A.; Zur, B. Adiabatic quantum transport in multiply connected systems. Rev. Modern Phys. 60 (1988), no. 4, 873–915

[BBK] Badanin, A; Brüning, J; Korotyaev, E. The Lyapunov function for Schrödinger operator with periodic 2×2 matrix potential, J. Funct. Anal. 234(2006), 106-126

[BBKL] Badanin, A; Brüning, J; Korotyaev, E.; Lobanov, I. Schrödinger operators on armchair graphs, in preparation

[BK] Badanin, A; Korotyaev, E. Spectral asymptotics for periodic forth order operators, Int. Math. Res. Not. 45(2005), 2775-2814

[Ca] Carlson, R. Spectral theory and spectral gaps for periodic Schrodinger operators on product graphs. Special section on quantum graphs. Waves Random Media 14 (2004), no. 1, S29–S45

[Ca1] Carlson, Robert Hill's equation for a homogeneous tree. Electron. J. Differential Equations 1997, No. 23, 30 pp. (electronic)

[CK] D. Chelkak; E, Korotyaev Spectral estimates for Schrödinger operator with periodic matrix potentials on the real line, Int. Math. Res. Not. 2006(2006), 0-41

[GT] Garnett J., Trubowitz E.: Gaps and bands of one dimensional periodic Schrödinger operators. Comment. Math. Helv. 59, 258-312 (1984)

[Ha] Harris P. Carbon Nanotubes and Related Structures, Cambridge Univ. Press., Cambridge, 1999

[KK] Kargaev P.; Korotyaev E. The inverse problem for the Hill operator, a direct approach. Invent. Math. 129, no. 3, 567-593 (1997), 138, no. 1, 227 (1999).

[KK1] Kargaev, P.; Korotyaev, E. Effective masses and conformal mappings. Comm. Math. Phys. 169 (1995), no. 3, 597–625

[Ko] Korotyaev, E. The inverse problem for the Hill operator. I Internat. Math. Res. Notices, 3(1997), 113–125

[KL] Korotyaev, E.; Lobanov, I. Zigzag periodic graphs in constant magnetic field, spectral theory, preprint 2006

[Ku] P. Kuchment, Graph models for waves in thin structures, *Waves in Random Media*, **12** (2002), R1-R24

[MW] Magnus, W.; Winkler, S. Hill's equation. Dover Publications, Inc., New York, 1979

[MO] Marchenko V.; Ostrovski I. A characterization of the spectrum of the Hill operator. Math. USSR Sb. 26, 493-554 (1975).

[MV] Molchanov, S.; Vainberg, B. Slowing down of the wave packets in quantum graphs. Waves Random Complex Media 15 (2005), no. 1, 101–112

[Pa] L. Pauling, The diamagnetic anisotropy of aromatic molecules, *Journal of Chemical Physics*, **4** (1936), 673-677.

[PT] Pöschel P., Trubowitz E.: *Inverse Spectral Theory*. Boston: Academic Press, 1987.

[ReS] Reed, M., Simon, B.: Methods of Modern Mathematical Physics, Vol.IV, Analysis of operators, Academic Press, New York, 1978

[ReS1] Reed, M., Simon, B.: Methods of Modern Mathematical Physics, Vol.II, Fourier Analysis, Self-Adjointness, Academic Press, New York, 1975

[RS] K. Ruedenberg and C. W. Scherr, Free-electron network model for conjugated systems. I. Theory, *The Journal of Chemical Physics*, **21** (1953), 1565-1581.

[SDD] Saito, R.; Dresselhaus, G.; Dresselhaus, M. Physical properties of carbon nanotubes, Imperial College Press, 1998

[SA] Schenker, J.; Aizenman, M. The creation of spectral gaps by graph decoration. Lett. Math. Phys. 53 (2000), no. 3, 253–262

[Ti] E. Titchmarsh, The theory of functions. Sec. ed., Univ. Press, London, 1975

[YS] Yakubovich, V., Starzhinskii, V. Linear differential equations with periodic coefficients. 1, 2. Halsted Press [John Wiley & Sons] New York-Toronto, 1975. Vol. 1, Vol. 2